

U-duality Twists and Possible Phase Transitions in 2+1D Supergravity

Ori J. Ganor¹

Department of Physics

Jadwin Hall

Princeton University

Princeton, NJ 08544, USA

We study 2+1D toroidal compactifications of M-theory with twists in the U-duality group. These compactifications realize many symmetric-manifolds from the classification of 2+1D extended supergravity moduli-spaces. We then focus on the moduli-space $SU(2,1)/U(2)$ obtained by dimensional reduction of pure $N = 2$ supergravity in 3+1D. This space is realized with an explicit example. Assuming that there are no quantum corrections, we conjecture that the classical discrete duality group has to be augmented with an extra strong/weak coupling duality. This implies the existence of new phases of the theory in which the original 8 compactification radii are all fixed at the Planck scale.

November, 1998

¹ origa@puhep1.princeton.edu

1. Introduction

In 2+1D the only massless bosonic propagating degrees of freedom are scalars. The restrictions on the moduli spaces of supergravity theories in 2+1D have been classified six years ago in [1]. For higher than $\mathcal{N} = 4$ supersymmetries, these moduli spaces must be of the form $\Gamma \backslash G/K$, where G is an appropriate non-compact group, K is its maximal compact subgroup and Γ is a discrete subgroup of G . Compactifications of M-theory provide a concrete realization of such theories and Γ is identified with the group of dualities. Thus, M-theory on $\mathbf{T}^8$ realizes the $\mathcal{N} = 16$ $E_{8(8)}/SO(16)$ moduli space of [2] and $E_8(\mathbb{Z})$ is the U-duality group [3]. Similarly, M-theory on $K_3 \times \mathbf{T}^4$ realizes the $SO(24, 8)/(SO(24) \times SO(8))$ moduli space and $SO(24, 8, \mathbb{Z})$ is the group of dualities [4].

One of the exciting recent developments is the physical interpretation of certain singularities in the moduli-space [5-10]. In general, the assumption of a free theory at low-energies breaks down when the theory is at a singular point in moduli space. A theory with $\mathcal{N} = 8$ rigid supersymmetries in 2+1D can have a singular point of the form $\mathbb{R}^{8n}/\Gamma$, where Γ is the Weyl-group of a certain Lie-algebra $\hat{g}$ of rank n . The low-energy description is then a strongly interacting conformal field theory. It is defined as the IR limit of 2+1D Super-Yang-Mills with the Lie-algebra $\hat{g}$. The low-energy description of theories with $\mathcal{N} = 4$ rigid supersymmetry at singularities is more complicated [8-10]. For a single vector multiplet it has been classified in [11]. In 3+1D singularities of the moduli space can be connections with another phase [5]. In 2+1D, singular points in moduli space of the form $\mathbb{R}^4/\Gamma$ are always connections with another phase (in all known cases).

One of the main motivations for the present work is to explore new phases of 2+1D gravity. One direction towards this goal is to study the singularities of the moduli space of the theory. The simplest moduli space of an $\mathcal{N} = 4$ supergravity theory is obtained by dimensional reduction of pure $\mathcal{N} = 2$ supergravity from 3+1D down to 2+1D. The classical moduli space is the homogeneous space $\Gamma_{cl} \backslash SU(2, 1)/(SU(2) \times U(1))$ where Γ_{cl} is a discrete subgroup (this moduli-space was introduced in [12-14]). The classical moduli space cannot encode the existence of other strongly-coupled phases. However, we will see that the structure of the moduli space might allow a consistent extension of Γ_{cl} to a U-duality group Γ . The extra dualities relate weak coupling to strong coupling and cannot be seen classically. The moduli space $\Gamma \backslash SU(2, 1)/(SU(2) \times U(1))$ has several singularities of the form $\mathbb{R}^4/\Gamma_0$ where Γ_0 is a finite subgroup of Γ . A 2+1D supergravity theory with such a moduli space will most likely have another phase emanating from these points in moduli space. In the other phases, the original coupling is stabilized.

To make the discussion more concrete we will construct a particular compactification of M-theory which realizes the moduli space $SU(2, 1)/(SU(2) \times U(1))$ (at least classically).

In fact, using what we know so far about M-theory, we can realize many of the other moduli-spaces found in [1] and hope to study their discrete duality groups. The compactification that we will use is similar in spirit to the compactifications presented in [15-19] and is obtained in two stages as follows. We start with a 3+1D vacuum of M-theory with $\mathcal{N} = 8$ supersymmetry. We can think of it as M-theory on a certain $\mathbf{T}^7$, but we pick a very special $\mathbf{T}^7$ (and fluxes) which make it a fixed point of the U-duality group $E_7(\mathbb{Z})$. Since U-duality is a discrete gauge symmetry, we can now compactify down to 2+1D on $\mathbf{S}^1$ with a U-duality twist. For an appropriate choice of the twist, we can preserve the desired amount of supersymmetry. The cases in [1] with more than $\mathcal{N} = 4$ had $\mathcal{N} = 5, 6, 8, 9, 10, 12, 16$ supersymmetries. We will construct examples which realize the cases $\mathcal{N} = 10, 12$ and some new cases with $\mathcal{N} = 8$. Given a particular U-duality element $\tilde{u} \in E_7(\mathbb{Z})$, we can imagine two ways of using it to compactify M-theory down to 2+1D. One way is to mod out by $\tilde{u}$ in 3+1D and then compactify on a circle. The second way is to leave the 3+1D limit intact but compactify on a circle with a twist. The advantage of the second kind of compactification is that we can be sure that no extra moduli come from “twisted sectors.”

As this work was in final stages, I found out that similar ideas have been put forward in [20] (referred to in [21]).¹ These papers discuss the universal hyper-multiplet in 3+1D and the discrete group Γ was studied in [20]. Although the setting in the present paper is physically different, the mathematical details are probably equivalent.

Furthermore, recently a paper appeared on the net [22] that presents a very comprehensive study of the moduli spaces of theories obtained via U-duality twists. In [22] only $D \geq 4$ dimensions were studied, but some of the examples presented in sections (2-3) of this paper are only a special case of [22].

2. U-duality twists

The compactifications that we will study are a special case of [15-19] and are of the following form. Start with M-theory on $\mathbf{T}^d$ which has the moduli space of,

$$\mathcal{M}_d = E_d(\mathbb{Z}) \backslash E_{d(d)}(\mathbb{R}) / K_d,$$

where K_d is the maximal compact subgroup of the non-compact $E_{d(d)}(\mathbb{R})$. The U-duality group $\Gamma_d \equiv E_d(\mathbb{Z})$ is, in general, not a symmetry of the compactification since it maps one point in $E_{d(d)}(\mathbb{R})$ to another. If Γ_d were acting freely, the U-duality group would not have been a symmetry at all. It would merely be the first homotopy group of $\mathcal{M}_d$. However,

¹ I am grateful to N. Seiberg for pointing out both these references.

Γ_d is not acting freely and for a given $\tilde{u} \in E_d(\mathbb{Z})$, there can be points in $E_{d(d)}(\mathbb{R})/K_d$ which are invariant under $\tilde{u}$. For example, for 9+1D type-IIB compactifications, if λ is the complex dilaton and $\tilde{u}$ is the generator of S-duality which maps $\lambda \rightarrow -1/\lambda$, then $\lambda = i$ is a fixed point of $\tilde{u}$. At $\lambda = i$ the element $\tilde{u}$ is an actual symmetry which acts non-trivially on the states. Furthermore, it is a gauge symmetry because it can be related to a geometrical rotation after compactification on a further $\mathbf{S}^1$ and mapping type-IIB on $\mathbf{S}^1$ to M-theory on $\mathbf{T}^2$. This situation is true in general. An element $\tilde{u} \in \Gamma_d$ is a gauge symmetry when the moduli of M-theory on $\mathbf{T}^d$ are derived from a point in $E_{d(d)}(\mathbb{R})/K_d$ which is fixed under $\tilde{u}$.

Now suppose we take a point $p \in E_{d(d)}(\mathbb{R})/K_d$ which is invariant under $\tilde{u}$ and then take M-theory on $\mathbf{T}^d$ with the moduli corresponding to the point p . Because $\tilde{u}$ is a gauge symmetry, we can compactify on an extra $\mathbf{S}^1$ with a twist of $\tilde{u}$. This means that if $\mathbf{S}^1$ is very large, after completing a full revolution around $\mathbf{S}^1$ any local state will turn into its $\tilde{u}$ -dual.

Similarly, we can take a base $\mathbf{T}^{d'}$ and d' U-duality twists $\tilde{u}_1, \dots, \tilde{u}_{d'}$ that commute with each other, and then compactify (M-theory on $\mathbf{T}^d$) further on $\mathbf{T}^{d'}$ to get a compactification down to $(11 - d - d')$. The next task would be to determine how much supersymmetry is left by such a twist.

2.1. Supersymmetry

Part of the U-duality group is actually the geometrical $SL(d, \mathbb{Z})$ symmetry. If all the twists are in $SL(d)$ then the compactification is just M-theory on a $(d + d')$ -dimensional manifold and the holonomy group of the manifold, which is generated by $\tilde{u}_1, \dots, \tilde{u}_{d'} \in SL(d)$ will determine the amount of supersymmetry. Sometimes, the group generated by $\tilde{u}_1, \dots, \tilde{u}_{d'}$ is not in any $SL(d)$ subgroup of Γ_d , but after compactification on an auxiliary $\mathbf{S}^1$ (unrelated to the $\mathbf{S}^1$ in the discussion above) and after conjugation by an element of the $(d + 1)$ -dimensional U-duality group Γ_{d+1} , all elements $\tilde{u}_1, \dots, \tilde{u}_{d'}$ can be made to lie inside the geometrical $SL(d + 1)$. This statement is actually a special case of the technique of [17] used to find generalizations of F-theory. A simple example of this statement is the U-duality group $SL(5, \mathbb{Z})$ for M-theory on $\mathbf{T}^4$ which is not entirely geometrical but after compactification on another $\mathbf{S}^1$ it can be conjugated inside $SO(5, 5, \mathbb{Z})$ to become the geometrical $SL(5, \mathbb{Z})$ of M-theory on $\mathbf{T}^5$ (see [17] for further details).

Before we determine the amount of supersymmetry left by the twist in the general case, let us find the fixed points. For simplicity we will take $d' = 1$. A fixed point is characterized by elements,

$$g \in E_{d(d)}(\mathbb{R}), \quad \tilde{k} \in K_d,$$

such that

$$\tilde{u} \circ g = g \circ \tilde{k}.$$

To determine the unbroken supersymmetries we need to know how $\tilde{u}$ acts on the spinors. The spinors are in a 2^r -dimensional representation of K_d . For $d = 8$ they are in the $(\mathbf{16}, \mathbf{2})$ of $SO(16) \otimes SO(2, 1)$. For $d = 7$ they are in the $(\mathbf{8}, \mathbf{2}) + (\bar{\mathbf{8}}, \mathbf{2}')$ of $SU(8) \otimes SO(3, 1)$. For $d = 6$ they are in the $(\mathbf{8}, \mathbf{4})$ of $Sp(4) \otimes SO(4, 1)$ (with an extra reality condition since both the $\mathbf{4}$ of $SO(4, 1)$ and $\mathbf{8}$ of $Sp(4)$ are pseudo-real). Since $\tilde{u}$ is conjugate to $\tilde{k} \in K_d$ it is easy to check that the fraction of unbroken supersymmetry is $l/2^r$ where l is the dimension of the eigen-space of $\tilde{k} = g^{-1}\tilde{u}g$ (in the relevant representation of K_d) with eigenvalue 1. For $d = 7$ we have $K_d = SU(8)$. Thus, to preserve at least $\frac{1}{4}$ of the supersymmetry, $\tilde{k} = g^{-1}\tilde{u}g$ must be in an $SU(6)$ subgroup of $SU(8)$.

2.2. Counting moduli

Let us now specialize to the case $d + d' = 8$. The compactification is down to 2+1D and we want to count the number of remaining moduli. Let us denote by Greek letters $(\mu, \nu, \dots = 0, 1, 2)$ space-time directions, by small English letters $(a, b, \dots = 3, \dots, d' + 2)$ the directions of the base $\mathbf{T}^{d'}$ and by capital English letters $(A, B, \dots = d' + 3, \dots, 10)$ the local directions of the fiber $\mathbf{T}^d$. Also, g denotes the metric and C the 3-form. The counting proceeds as follows:

1. There are $\frac{d'(d'+1)}{2}$ moduli for the metric g_{ab} .
2. There are d' moduli for the duals of the vectors $g_{\mu a}$.
3. There are $\frac{d'(d'-1)(d'-2)}{6}$ moduli for C_{abc} .
4. There are $\frac{d'(d'-1)}{2}$ moduli for the duals of the vectors $C_{\mu ab}$.
5. We need to know n_s , the number of scalar moduli of M-theory on $\mathbf{T}^d$ that are preserved by the twists.
6. We need to know n_v , the number of vectors of M-theory on $\mathbf{T}^d$ that are preserved by the twists. We then get $(d' + 1)n_v$ moduli in 2+1D if $d' > 1$. The extra one is the dual of the vector in 2+1D. When $d = 7$ and $d' = 1$ we get just n_v and not $2n_v$, because n_v counts electric and magnetic duals twice.
7. For $d < 5$ we need to know how many tensors n_T of M-theory on $\mathbf{T}^d$ are preserved. We then obtain $\frac{d'(d'-1)}{2}n_T$ moduli of the form T_{ab} and $d'n_T$ moduli which are duals of $T_{a\mu}$. For $d = 5$ and $d' = 3$ we get only $3n_T$ because we count electric-magnetic duals twice.
8. We will not consider cases $d \leq 3$ here.

Now let us specialize to $d = 7$ and $d' = 1$. To count the surviving scalar moduli, n_s , we decompose E_7 under $SU(8) \subset E_7$. The 70 scalar moduli are in the irreducible

representation, **70**, of $SU(8)$. This representation is real and can be represented as the space of the anti-symmetric 4-forms. We need to know how many eigenvalues of 1 the element $\tilde{k} = g^{-1}\tilde{u}g$ possesses in this representation. To count the vectors n_v we need to know how many eigenvalues of 1 appear among the eigenvalues of $\tilde{k}$ in the representation, **28** + $\overline{\mathbf{28}}$, of the vectors.

In the general case of modding out by a U-duality twist, as studied in [18], one cannot tell, at present, whether there are more moduli that cannot be obtained by simply reducing the spectrum of M-theory on $\mathbf{T}^d$. For example, if we counted the number of vectors of M-theory on $\mathbf{T}^4/\mathbb{Z}_2$ by reducing the spectrum of M-theory on $\mathbf{T}^4$ we would not get the 16 blow-up modes. This $\mathbb{Z}_2$ was a subgroup of $SL(5, \mathbb{Z})$ and it is hard to tell what happens for twists which do not even have a weakly coupled string-theory limit. On the other hand, in the special cases that we study, one can be sure that there are no new moduli, since for a large $\mathbf{T}^{d'}$ there is no singularity and the classical approximation is good. As an example, if we compactified M-theory on $\mathbf{T}^4$ further down on $\mathbf{S}^1$ with the same $\mathbb{Z}_2$ twist as above, we would have got a smooth 5-manifold with $SU(2)$ holonomy. It gives a 5+1D vacuum with $\mathcal{N} = (1, 1)$ supersymmetry and only 8 vectors with an,

$$SO(4, 4, \mathbb{Z}) \backslash SO(4, 4, \mathbb{R}) / (SO(4) \times SO(4)),$$

moduli space. This is in contrast to compactifying M-theory on $\mathbf{T}^4/\mathbb{Z}_2$ further down on $\mathbf{S}^1$ with the,

$$SO(20, 4, \mathbb{Z}) \backslash SO(20, 4, \mathbb{R}) / (SO(20) \times SO(4)),$$

moduli space.

3. Examples

We now turn to a few examples. We will start with the cases in 5+1D and then proceed to 2+1D. Note that only the cases with $\mathcal{N} = 8, 16$ in 2+1D can conceivably be (untwisted) toroidal compactifications of a 3+1D model.

3.1. Examples in 5+1D

We start with M-theory on $\mathbf{T}^4$ and compactify down on $\mathbf{S}^1$ with a U-duality twist that preserves half the supersymmetry. The U-duality twist is conjugate to an $Sp(2)$ element. Any element in $Sp(2)$ is conjugate to an element in an $SO(4)$ subgroup. Thus the element $\tilde{u}$ is conjugate to a geometrical twist. This does not necessarily mean that all twists are

geometrical since there might be two elements $\tilde{u}_1$ and $\tilde{u}_2$ which are conjugate in $SL(5, \mathbb{R})$ but non-conjugate in $SL(5, \mathbb{Z})$.

The twist can now be represented as $e^{i\alpha} \in U(1) \subset SU(2) \subset SO(4)$. The particular embedding is such that the fundamental $\mathbf{4}$ of $SO(4)$ becomes two spinors of $SU(2)$. The possible values of α are,

$$\alpha = \frac{\pi}{3}, \frac{\pi}{2}, \frac{2\pi}{3}.$$

The possible geometrical twists are thus as follows (up to conjugacy in $SL(4, \mathbb{R})$):

1. We can twist by the $\mathbb{Z}_2$ acting on all 4 directions of a generic $\mathbf{T}^4$.
2. We take a special $\mathbf{T}^4$ which is a product $\mathbf{T}^2 \times \mathbf{T}^2$ and each $\mathbf{T}^2$ has a complex structure which is fixed to one of the values

$$\tau = e^{\frac{\pi i}{2}}, e^{\frac{\pi i}{3}}.$$

We determine the number of vectors in 5+1D as follows. Let $I, J, K, \dots$ denote directions inside $\mathbf{T}^4$ and let 6 denote the direction of the extra $\mathbf{S}^1$. $\mu, \nu, \dots$ denote 5+1D space-time directions. The dual of the 3-form of M-theory and the graviton field $g_{\mu 6}$ give rise to two vectors in 5+1D. The vectors of the form $C_{\mu 6 I}$ and the form $g_{\mu I}$ do not survive the $\tilde{u}$ -twist. The number of vectors of the form $C_{\mu I J}$ which survive the $\tilde{u}$ -twist is equal to the dimension of the invariant subspace of $\tilde{u}$ in $\mathbf{6}$ of $SO(4)$. This number is 6 if we have a $\mathbb{Z}_2$ twist, and is 4 for the other cases. The moduli space is,

$$\Gamma \backslash SO(4, 4, \mathbb{R}) / (SO(4) \times SO(4)),$$

in the case of the $\mathbb{Z}_2$ twist, and is,

$$\Gamma \backslash SO(4, 2, \mathbb{R}) / (SO(4) \times SO(2)),$$

in the case of the special twists. Compactifying these models to 2+1D gives examples of theories with moduli spaces,

$$\Gamma \backslash SO(8, 8, \mathbb{R}) / (SO(8) \times SO(8)),$$

and,

$$\Gamma \backslash SO(8, 6, \mathbb{R}) / (SO(8) \times SO(6)),$$

respectively.

3.2. Examples in 2+1D

We now turn to a few examples in 2+1D. We start with M-theory on $\mathbf{T}^7$ and compactify on $\mathbf{S}^1$ with a twist $\tilde{u} \in E_7(\mathbb{Z})$. We saw in the previous section that to preserve $\mathcal{N} = 2r$ supersymmetries in 2+1D, we need a twist $\tilde{u}$ which is conjugate in $E_{7(7)}(\mathbb{R})$ to an element $\tilde{k} \in SU(8)$ with exactly r eigenvalues of 1. In the classification of [1] there are also cases with $\mathcal{N} = 9$ and $\mathcal{N} = 5$. They cannot be realized by the kind of compactifications we are studying. Returning to $\mathcal{N} = 2r$, we will calculate the number of moduli $k = 2 + n_v + n_s$ in each case. Two moduli are the radius Λ of the circle and the 2+1D dual Φ of the graviton $g_{\mu 3}$ where 3 is the direction of the circle. n_v is the number of 3+1D vectors of the compactification of M-theory on $\mathbf{T}^7$ which are left invariant by $\tilde{u}$ and n_s is the number of scalars which are left invariant.

It is also interesting to check whether the twist can be realized geometrically or not. If the twist is realized geometrically $\tilde{u}$ must be in an $SO(7) \subset SU(8)$ subgroup. The embedding of $SO(7)$ is such that the fundamental $\mathbf{8}$ of $SU(8)$ becomes the spinor $\mathbf{8}$ of $SO(7)$. We can embed it as $SO(7) \subset SO(8) \subset SU(8)$ where the first embedding is such that the vector $\mathbf{8}_v$ of $SO(8)$ becomes $\mathbf{7} + \mathbf{1}$ of $SO(7)$ and the second embedding is such that $\mathbf{8}$ of $SU(8)$ becomes the spinor $\mathbf{8}_s$ of $SO(8)$. Given an element $g \in SO(8)$, its matrix in the vector representation $\mathbf{8}_v$ can be brought into a block diagonal form with four 2×2 blocks of the form,

$$\begin{pmatrix} \cos \alpha_i & \sin \alpha_i \\ -\sin \alpha_i & \cos \alpha_i \end{pmatrix}, \quad -\pi \leq \alpha_i \leq \pi.$$

The quadruple $(\alpha_1, \alpha_2, \alpha_3, \alpha_4)$ is determined up to permutation and up to a change in sign,

$$(\alpha_1, \alpha_2, \alpha_3, \alpha_4) \longrightarrow (\epsilon_1 \alpha_1, \epsilon_2 \alpha_2, \epsilon_3 \alpha_3, \epsilon_4 \alpha_4) \quad \epsilon_i = \pm 1, \quad \prod \epsilon_i = 1.$$

The matrix of g in the spinor representation $\mathbf{8}_s$ has a similar representation with a quadruple $(\beta_1, \dots, \beta_4)$ where

$$\begin{aligned} \beta_1 &= \frac{\alpha_1 + \alpha_2 + \alpha_3 + \alpha_4}{2}, \\ \beta_2 &= \frac{\alpha_1 + \alpha_2 - \alpha_3 - \alpha_4}{2}, \\ \beta_3 &= \frac{\alpha_1 - \alpha_2 - \alpha_3 + \alpha_4}{2}, \\ \beta_4 &= \frac{\alpha_1 + \alpha_2 - \alpha_3 - \alpha_4}{2}. \end{aligned} \tag{3.1}$$

(modulo 2π). There is of course the ambiguity of π in each phase. Now, if an element of $SU(8)$ is really an element of $SO(8)$ then all the eigenvalues of the matrix which represents

it in **8** of $SU(8)$ must come in pairs of $e^{\pm i\alpha_i}$ for some 4 phases α_i . In addition, if it is an element of $SO(7)$ in the above embedding then it must be that one of the β_i 's is 0. Thus, the condition that a twist $\tilde{u}$ is geometrical is that the eigenvalues of $\tilde{u}$, as an $SU(8)$ matrix, are of the form (for a particular choice of labeling of the eigenvalues),

$$e^{\pm i\alpha_i}, \quad i = 1 \dots 4, \quad e^{i \sum_1^4 \alpha_i} = 1.$$

It is also interesting to check if $\tilde{u}$ can be represented as a T-duality element of type-IIA or type-IIB on $\mathbf{T}^6$. In this case $\tilde{u}$ must be in the,

$$SO(6) \otimes SO(6) \sim SU(4) \otimes SU(4) \subset SU(8),$$

subgroup of $SU(8)$ which means that it must be possible to group the 8 eigenvalues in two groups of 4 such that the product in each group is 1.

Finally, in each case listed below, we will identify the eigenvalues of $\tilde{u}$ as an $SU(8)$ matrix. We will not attempt to find a $g \in E_{7(7)}(\mathbb{R})$ such that $gug^{-1} \in E_7(\mathbb{Z})$ but we will check that the characteristic polynomial $\det(xI - \tilde{u})$ in the representation, **28** + $\overline{\mathbf{28}}$, of $SU(8)$ has integer coefficients.

Cases with $\mathcal{N} = 12$ supersymmetry:

$\tilde{u} \in SU(8)$ must have 6 eigenvalues of 1. The other two eigenvalues of $\tilde{k}$ must be $e^{\pm i\theta}$, and $\tilde{k} \in U(1) \subset SU(8)$. The possible values of θ are determined by the requirement that in the representation **56** of $SU(8)$, the characteristic polynomial, $\det(xI - \tilde{k})$, should be integral. It is easy to calculate,

$$\det_{\mathbf{56}}(xI - \tilde{k}) = (x - 1)^{32}(x - e^{i\theta})^{12}(x - e^{-i\theta})^{12}.$$

This leaves the possibilities,

$$\theta = \frac{\pi}{3}, \frac{\pi}{2}, \frac{2\pi}{3}, \pi.$$

The number of vectors n_v is determined by decomposing **28** + $\overline{\mathbf{28}}$ of $SU(8)$ under $U(1) \subset SU(8)$,

$$\mathbf{28} + \overline{\mathbf{28}} = 12(\mathbf{1}_{-1}) + 12(\mathbf{1}_{+1}) + 32(\mathbf{1}_0).$$

which leaves us with $n_v = 32$. This number counts a vector and its dual separately. As for n_s , we need to decompose the representation **70** of $SU(8)$ under $U(1) \subset SU(8)$. We find that,

$$\mathbf{70} = 30(\mathbf{1}_0) + 20(\mathbf{1}_{+1}) + 20(\mathbf{1}_{-1}).$$

This leaves us with $n_s = 30$. Altogether, the moduli space has dimension,

$$2 + n_s + n_v = 64,$$

which agrees with the entry in table of [1] for $\mathcal{N} = 12$ (repeated in appendix A).

According to the discussion at the beginning, $\tilde{u}$ is not a geometrical twist, but can be put in the form of an $SO(6, 6, \mathbb{Z})$ T-duality twist.

Cases with $\mathcal{N} = 10$ supersymmetry:

This time $\tilde{u} \in SU(8)$ has to have exactly 5 eigenvalues of 1. Let the other eigenvalues be

$$e^{i\alpha_1}, e^{i\alpha_2}, e^{i\alpha_3}, \quad \alpha_1 + \alpha_2 + \alpha_3 = 0.$$

The characteristic polynomial in $\mathbf{28} + \overline{\mathbf{28}}$ of $SU(8)$ is,

$$\det_{\mathbf{56}}(xI - \tilde{k}) = (x - 1)^{20} P(x)^6, \quad (3.2)$$

$$P(x) \equiv (x - e^{i\alpha_1})(x - e^{-i\alpha_1})(x - e^{i\alpha_2})(x - e^{-i\alpha_2})(x - e^{i\alpha_3})(x - e^{-i\alpha_3})$$

This polynomial has integral coefficients only in the the following cases, listed up to an S_3 permutation and up to an overall minus sign (replacing $\tilde{k}$ by $\tilde{k}^{-1}$ which clearly does not make any difference)

$$\left(\frac{\alpha_1}{2\pi}, \frac{\alpha_2}{2\pi}, \frac{\alpha_3}{2\pi}\right) = \left(\frac{1}{7}, \frac{2}{7}, \frac{4}{7}\right), \left(\frac{1}{6}, \frac{1}{6}, \frac{4}{6}\right), \left(\frac{1}{6}, \frac{2}{6}, \frac{3}{6}\right), \left(\frac{1}{4}, \frac{1}{4}, \frac{2}{4}\right), \left(\frac{1}{3}, \frac{1}{3}, \frac{1}{3}\right). \quad (3.3)$$

Since $\tilde{u} \in SU(3)$, to determine n_v and n_s we have to decompose the appropriate representations of $SU(8)$ under $SU(3) \subset SU(8)$.

$$\begin{aligned} \mathbf{28} + \overline{\mathbf{28}} &= 20(\mathbf{1}) + 6(\mathbf{3}) + 6(\overline{\mathbf{3}}), \\ \mathbf{70} &= 10(\mathbf{1}) + 10(\mathbf{3}) + 10(\overline{\mathbf{3}}). \end{aligned} \quad (3.4)$$

Thus, in all cases above, $n_v = 20$ and $n_s = 10$. Altogether, $n_v + n_s + 2 = 32$ as expected from the table in appendix (A).

Again, all of these twists are not geometrical but can be represented as $SO(6, 6, \mathbb{Z})$ T-duality twists.

Cases with $\mathcal{N} = 8$ supersymmetry:

We denote the four eigenvalues of $\tilde{k}$ which are not 1 by,

$$e^{i\alpha_j}, \quad j = 1 \dots 4, \quad \sum \alpha_j = 0 \pmod{4}.$$

The characteristic polynomial in $\mathbf{28} + \overline{\mathbf{28}}$ of $SU(8)$ is,

$$\begin{aligned} \det_{\mathbf{56}}(xI - \tilde{k}) &= (x-1)^{12} P(x)^4 Q(x)^{12}, \\ P(x) &\equiv \prod_1^4 (x - e^{i\alpha_j}) \prod_1^4 (x - e^{-i\alpha_j}), \\ Q(x) &\equiv \prod_{1 \leq i < j \leq 4} (x - e^{i(\alpha_i + \alpha_j)}) \end{aligned} \quad (3.5)$$

It has integral coefficients for,

$$n = 2, 3, 4, 5, 6, 7, 8, 9, 10, 12, 14, 15, 16, 18, 20, 24, 30.$$

$$\begin{aligned} \left(\frac{\alpha_1}{2\pi}, \frac{\alpha_2}{2\pi}, \frac{\alpha_3}{2\pi}, \frac{\alpha_4}{2\pi}\right) &= \left(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}\right), \left(\frac{1}{3}, \frac{1}{3}, \frac{2}{3}, \frac{2}{3}\right), \left(\frac{1}{4}, \frac{1}{4}, \frac{1}{4}, \frac{1}{4}\right), \left(\frac{1}{4}, \frac{1}{4}, \frac{3}{4}, \frac{3}{4}\right), \left(\frac{1}{4}, \frac{3}{4}, \frac{2}{4}, \frac{2}{4}\right), \\ &\left(\frac{1}{5}, \frac{2}{5}, \frac{3}{5}, \frac{4}{5}\right), \left(\frac{1}{6}, \frac{1}{6}, \frac{1}{6}, \frac{3}{6}\right), \left(\frac{1}{6}, \frac{1}{6}, \frac{2}{6}, \frac{2}{6}\right), \left(\frac{1}{6}, \frac{1}{6}, \frac{5}{6}, \frac{5}{6}\right), \left(\frac{1}{6}, \frac{2}{6}, \frac{4}{6}, \frac{5}{6}\right), \\ &\left(\frac{1}{6}, \frac{3}{6}, \frac{3}{6}, \frac{5}{6}\right), \left(\frac{1}{8}, \frac{1}{8}, \frac{3}{8}, \frac{3}{8}\right), \left(\frac{1}{8}, \frac{3}{8}, \frac{5}{8}, \frac{7}{8}\right), \left(\frac{1}{9}, \frac{2}{9}, \frac{3}{9}, \frac{4}{9}\right), \left(\frac{1}{10}, \frac{3}{10}, \frac{7}{10}, \frac{9}{10}\right), \\ &\left(\frac{1}{12}, \frac{5}{12}, \frac{7}{12}, \frac{11}{12}\right), \left(\frac{2}{12}, \frac{3}{12}, \frac{9}{12}, \frac{10}{12}\right), \left(\frac{1}{15}, \frac{2}{15}, \frac{4}{15}, \frac{8}{15}\right), \left(\frac{1}{15}, \frac{2}{15}, \frac{4}{15}, \frac{8}{15}\right), \\ &\left(\frac{1}{20}, \frac{9}{20}, \frac{13}{20}, \frac{17}{20}\right), \left(\frac{1}{20}, \frac{3}{20}, \frac{7}{20}, \frac{9}{20}\right), \left(\frac{1}{24}, \frac{5}{24}, \frac{7}{24}, \frac{11}{24}\right), \\ &\left(\frac{1}{24}, \frac{11}{24}, \frac{17}{24}, \frac{19}{24}\right), \left(\frac{1}{30}, \frac{17}{30}, \frac{19}{30}, \frac{23}{30}\right). \end{aligned} \quad (3.6)$$

To calculate n_s and n_v we decompose the appropriate representations of $SU(8)$ under $SU(4) \subset SU(8)$.

$$\begin{aligned} \mathbf{28} + \overline{\mathbf{28}} &= 12(\mathbf{1}) + 4(\mathbf{4}) + 4(\overline{\mathbf{4}}) + 2(\mathbf{6}), \\ \mathbf{70} &= 2(\mathbf{1}) + 4(\mathbf{4}) + 4(\overline{\mathbf{4}}) + 6(\mathbf{6}). \end{aligned} \quad (3.7)$$

In all the cases above, there is no vector of $\mathbf{4}$ or $\overline{\mathbf{4}}$ which is left invariant. Let $2l$ be the number of vectors of $\mathbf{6}$ which are left invariant under $\tilde{u}$. This is given by the number of pairs of eigenvalues which sum up to zero. In the cases above there are examples with $l = 0, 1, 2, 3$. Then,

$$2 + n_s + n_v = 2 + 2 + 12 + 16l = 16 + 16l.$$

Thus, the moduli space is $SO(8, k)/(SO(8) \otimes SO(k))$ with $k = 2, 4, 6, 8$. Note that this time some of the $\tilde{u}$ listed above can be represented as purely geometrical twists. In fact, all the geometrical $\tilde{u}$'s can be represented as elements of $SU(6)$ which means that they

can be realized as ordinary compactifications of 3+1D models with $\mathcal{N} = 4$ supersymmetry on $\mathbf{S}^1$. This does not mean that all *vacua* can be represented in this way. In principle, there could still be inequivalent elements $g \in E_{7(7)}(\mathbb{R})$ such that $g^{-1}\tilde{u}g \in E_7(\mathbb{Z})$ are not conjugate in $E_7(\mathbb{Z})$ (although they are conjugate in $E_{7(7)}(\mathbb{R})$).

There are a few cases which are toroidal compactification of a 5+1D model with $\mathcal{N} = (1, 1)$ supersymmetry. This can be obtained by compactifying M-theory on $\mathbf{T}^4$ further down on $\mathbf{S}^1$ with a U-duality twist which preserves half the supersymmetry. These are the cases discussed in the first subsection.

Cases with $\mathcal{N} = 6$ supersymmetry:

In this case we are looking for 5 eigenvalues,

$$e^{i\alpha_i}, \quad \sum_1^5 \alpha_i = 0 \pmod{2\pi}.$$

We will not attempt to exhaust all the cases. We will just point out that, unlike the previous cases, for $\mathcal{N} = 6$ there are examples where $\tilde{u}$ cannot be conjugated to a T-duality $SO(6, 6, \mathbb{Z})$ element. For example,

$$\left(\frac{\alpha_1}{2\pi}, \frac{\alpha_2}{2\pi}, \frac{\alpha_3}{2\pi}, \frac{\alpha_4}{2\pi}, \frac{\alpha_5}{2\pi}\right) = \left(\frac{1}{11}, \frac{3}{11}, \frac{4}{11}, \frac{5}{11}, \frac{9}{11}\right), \quad (3.8)$$

We confess to not having checked that there really is a U-duality element in $E_{7(7)}(\mathbb{R})$ which conjugates this $SU(8)$ element into an $E_7(\mathbb{Z})$ element. All we know is that its characteristic polynomial is integral.

3.3. Examples with $\mathcal{N} = 4$ in 2+1D

We now present a few examples with $\mathcal{N} = 4$.

12 moduli:

Here is a variant of a model that has appeared several times in the literature. Take a base $\mathbf{T}^2$ and a fiber of the form $\mathbf{T}^6 = (\mathbf{T}^2)^3$ such that each $\mathbf{T}^2$ has $\tau = \frac{1}{2} + \frac{\sqrt{3}}{2}i$, i.e. forms a hexagonal lattice. We also require all the $\mathbf{T}^2$'s to have equal area. Let z_1, z_2, z_3 be complex coordinates on the three $\mathbf{T}^2$'s, with the identifications,

$$z_i \sim z_i + 1 \sim z_i + \frac{1}{2} + \frac{\sqrt{3}}{2}i.$$

Let us define the elements $\tilde{u}_1, \tilde{u}_2 \in SU(3)$ as follows.

$$\begin{aligned}\tilde{u}_1 : (z_1, z_2, z_3) &\rightarrow (\omega z_1, \omega z_2, \omega z_3), \\ \tilde{u}_2 : (z_1, z_2, z_3) &\rightarrow (z_3, z_1, z_2),\end{aligned}\tag{3.9}$$

where $\omega = e^{\frac{i\pi}{3}}$. $\tilde{u}_1$ and $\tilde{u}_2$ preserve the complex structure and the holomorphic 3-form. Now we compactify on $\mathbf{T}^2$ with the twists $\tilde{u}_1$ and $\tilde{u}_2$ in the fiber $\mathbf{T}^6$ as we go along the sides of $\mathbf{T}^2$. The resulting manifold is a 4-fold with $SU(3)$ holonomy. Compactification of M-theory on this manifold preserves $\frac{1}{4}$ -supersymmetry. The remaining moduli are listed below. We denote by i, j, k holomorphic coordinates and by $\bar{i}, \bar{j}, \dots$ anti-holomorphic coordinates. a, b denote directions on the base and $\mu, \nu = 0, 1, 2$ are space-time directions. C_{ABC} denotes the VEV of the M-theory 3-form. The moduli are:

1. The shape and size of the base $\mathbf{T}^2$ as well as the dual of the 1-form $C_{ab\mu}$ made out of the 3-form of M-theory integrated on $\mathbf{T}^2$. Altogether this gives 4 moduli.
2. The volume of $\mathbf{T}^6$ which is one modulus.
3. The 3-form integrated on the $(3, 0)$ and $(0, 3)$ cycles, i.e. C_{ijk} and $C_{i\bar{j}\bar{k}}$. Altogether 2 moduli.
4. The moduli of the form $C_{ij\bar{k}}$ and $C_{i\bar{j}\bar{k}}$ get multiplied by ω and ω^{-1} respectively under the operation of $\tilde{u}_1$ and therefore become massive.
5. The only moduli of the form $C_{i\bar{j}a}$ which are invariant under $\tilde{u}_1$ and $\tilde{u}_2$ are,

$$\sum_{k=1}^3 C_{k\bar{k}a}.$$

This gives 2 more moduli. We also need to add the dual of $\sum_{k=1}^3 C_{k\bar{k}\mu}$ to obtain a third modulus.

6. The duals of the gravitons $g_{\mu a}$ give two more moduli. Altogether we have 12 moduli.

20 moduli:

We can pick a particular $\mathbf{T}^6$ and an element $\tilde{u} \in SU(3)$ such that $\tilde{u}^7 = I$ and such that $\tilde{u}$ preserves only the volume of $\mathbf{T}^6$. To describe $\tilde{u}$, we start with $\mathbb{R}^7$ with coordinates $(x_1, x_2, \dots, x_7)$ and take $\tilde{u}_0 \in \mathbb{Z}_7 \subset SO(7)$ to be the cyclic permutation. Since $\tilde{u}_0$ preserves the diagonal direction it is actually in $SO(6)$. Let $\tilde{u}$ be the reduced action of $\tilde{u}_0$ on the space $\mathbb{R}^6$ that is orthogonal to the diagonal $(1, 1, 1, 1, 1, 1)$. Its eigenvalues are $e^{\frac{2\pi i k}{7}}$ for $k = 1, \dots, 6$. In the spinor representation $\mathbf{4}$ of $SO(6)$, its eigenvalues are $e^{\frac{2\pi i}{7}}, e^{\frac{4\pi i}{7}}, e^{\frac{8\pi i}{7}}$ and 1. It thus preserves one spinor. To describe $\mathbf{T}^6$, take the 7 points in $\mathbb{R}^7$,

$$(1, 0, 0, 0, 0, 0, 0), (0, 1, 0, 0, 0, 0, 0), \dots, (0, 0, 0, 0, 0, 0, 1).$$

They form a 6-plex which is $\tilde{u}_0$ invariant. Taking one of these as a base, the vectors to the other six are unit vectors of a lattice in $\mathbb{R}^6$. Its unit cell is the requisite $\mathbf{T}^6$. Now, compactify on a further $\mathbf{S}^1$ with a $\tilde{u}$ twist. It is easy to check that we end up with a 3+1D model with $\mathcal{N} = 2$ supersymmetry. The low-energy description has 3 vector multiplets, 1 hyper-multiplet and a gravity multiplet (containing a graviphoton). The radius of the circle is in a hyper-multiplet so the metric on the vector-multiplet moduli space can be calculated for large R and does not get quantum corrections [23]. It would be interesting to find a heterotic dual to this compactification, perhaps along the lines of [15]. After further compactification to 2+1D we find 20 scalar moduli and $k = 5$.

4 moduli:

We now turn to an example which involves a non-geometrical twist and has only 4 moduli. This is the example that we will study in detail later on. We start with M-theory on $\mathbf{T}^7$ and compactify on $\mathbf{S}^1$ with a twist $\tilde{u} \in E_7(\mathbb{Z})$. We saw in the previous section that to preserve 8 supersymmetries, i.e. $\mathcal{N} = 4$ in 2+1D, we need a twist $\tilde{u}$ which is conjugate in $E_{7(7)}(\mathbb{R})$ to an element $\tilde{k} \in SU(6) \subset SU(8)$. In this case the number of moduli will be $k = 2 + n_v + n_s$. Two moduli are the radius Λ of the circle and the 2+1D dual Φ of the graviton $g_{\mu 3}$ where “3” is the direction of the circle. n_v is the number of 3+1D vectors of the compactification of M-theory on $\mathbf{T}^7$ which are left invariant by $\tilde{u}$ and n_s is the number of scalars which are left invariant.

We now choose $\tilde{k}$ to be the diagonal matrix with eigenvalues,

$$\underbrace{(e^{\frac{i\pi}{3}}, \dots, e^{\frac{i\pi}{3}})}_{6 \text{ times}}, 1, 1). \quad (3.10)$$

In this way, $\tilde{k}$ is in the center $\mathbb{Z}_6 \subset SU(6)$. The number of vectors n_v is determined by decomposing **28** of $SU(8)$ under $SU(6)$,

$$\mathbf{28} = \mathbf{15} + 2(\mathbf{6}) + \mathbf{1}.$$

Under the $\mathbb{Z}_6$ center, only **1** is invariant which gives us (together with $\overline{\mathbf{28}}$) $n_v = 2$. The two vectors are electric-magnetic duals. As for n_s , we need to decompose the representation **70** of $SU(8)$ under $SU(6) \subset SU(8)$. We find that,

$$\mathbf{70} = \mathbf{15} + \overline{\mathbf{15}} + \mathbf{40}.$$

here **40** is made of the anti-symmetric 3-forms and is real. These representations are all charged under $\mathbb{Z}_6$ and therefore $n_s = 0$.

The eigenvalues of $\tilde{u}$ cannot be separated into two groups whose product is 1. Therefore, the twist is not conjugate to a T-duality twist. This is also obvious from the fact that there is no extra modulus which could be a string coupling constant.

It remains to prove that there actually exists an element $\tilde{u} \in E_7(\mathbb{Z})$ which is conjugate to $\tilde{k}$ above. Let us first note the following. Denote,

$$\omega = e^{\frac{i\pi}{3}}, \quad \omega^6 = 1.$$

In the representation **56**, the eigenvalues of $\tilde{k}$ are as follows. ω appears 12 times, ω^{-1} appears 12 times, ω^2 appears 15 times, ω^{-2} appears 15 times and 1 appears twice. The characteristic polynomial is thus,

$$\begin{aligned} \det(xI - \tilde{k}) &= (x - \omega)^{12}(x - \omega^{-1})^{12}(x - \omega^2)^{15}(x - \omega^{-2})^{15}(x - 1)^2 \\ &= (x^2 - x + 1)^{12}(x^2 + x + 1)^{15}(x - 1)^2. \end{aligned} \tag{3.11}$$

It has integral coefficients, as a matrix that is conjugate to a matrix in $E_7(\mathbb{Z})$ should have. Note also that the trace is (-1) which means that $\tilde{u}$ cannot be separated into two blocks of 28×28 and it mixes both electric and magnetic charges.

In appendix (C), we will construct an example of such a $\tilde{u}$ explicitly.

4. The classical limit

In the remaining of this paper we will concentrate on the last example from the previous section. This example was the compactification of M-theory on a fixed $\mathbf{T}^7$ further down on $\mathbf{S}^1$ of radius Λ with a U-duality twist that fixes the moduli of $\mathbf{T}^7$. Most of the time, the particular details of the twist will not concern us. The moduli space is a quaternionic manifold of dimension 4. We will assume below that it is $\Gamma \backslash SU(2, 1) / (SU(2) \times U(1))$. Locally, this is the same moduli space that one obtains from dimensional reduction of $\mathcal{N} = 2$ 3+1D supergravity (see [12-14]). To see why we could get the same moduli space in our case we will first study the classical limit of large radius Λ . In this limit, we can first take the low-energy limit of M-theory on $\mathbf{T}^7$ at the particular fixed moduli, and then compactify that 3+1D low-energy action further down on $\mathbf{S}^1$ with a twist. This process involves a subtlety which we shall now discuss.

4.1. *S-duality twists*

The low-energy of M-theory on $\mathbf{T}^7$ has 56 electric and magnetic field-strengths. We have seen that the twist along $\mathbf{S}^1$ is not a mere geometrical transformation (otherwise, the volume of $\mathbf{T}^7$ would remain a modulus) and it takes some of the charges to their magnetic duals. How do we twist an abelian $U(1)$ theory by S-duality or, more generally, by the $SL(2, \mathbb{Z})$ group? This topic was discussed in [24] in the context of the D3-brane partition function in F-theory. The result is as follows. Let $0 \leq \xi \leq 2\pi$ be the coordinate on $\mathbf{S}^1$. To describe the twist, we must allow the gauge field $A_\mu(x, \xi)$ to be discontinuous at $\xi = 0$. We take the action,

$$\frac{1}{4g^2} \int d\xi d^3x F^2,$$

in the bulk, and add to it a 2+1D Chern-Simons-like interaction,

$$\frac{1}{2c} \int d^3x [aA \wedge dA + 2A \wedge dA' - fA' \wedge dA'].$$

Here,

$$A_\mu(x) \equiv A_\mu(x, 0), \quad A'_\mu(x) \equiv A_\mu(x, 2\pi),$$

and $\begin{pmatrix} a & b \\ c & f \end{pmatrix}$ is the $SL(2, \mathbb{Z})$ duality matrix. In general, the extra Chern-Simons interaction will make the gauge field massive. The only massless gauge fields which remain are those which are invariant under the twist.

4.2. *Classical Kaluza-Klein reduction*

The action of the remaining zero-modes in 2+1D is determined, in the limit $\Lambda \rightarrow \infty$, by dimensionally reducing the classical action of 3+1D. The 3+1D action contains 70 scalar fields which, after the twist, all become massive. It also contains 56 $U(1)$ gauge-fields and their magnetic duals and we have seen that only 2 of them are invariant under the $\tilde{u}$ -twist. These two must therefore be electric-magnetic duals of each other. To get the 2+1D dimensionally reduced action, it is thus enough to start with a 3+1D action of the form,

$$\int d^3x d\xi \sqrt{g} [R(g) + \frac{1}{4e^2} \tilde{H}_{\mu\nu} \tilde{H}^{\mu\nu} + \frac{\theta}{32\pi^2} \epsilon^{\mu\nu\beta\gamma} \tilde{H}_{\mu\nu} \tilde{H}_{\beta\gamma}].$$

Here ξ is the coordinate along $\mathbf{S}^1$, $g_{\mu\nu}$ is the 3+1D metric, $\tilde{H}$ is the field-strength of the $U(1)$ gauge field $\tilde{B}_\mu$ and the coupling constant is e . The coupling constant can be determined from the fixed point g of $\tilde{u}$ in the moduli space of M-theory on $\mathbf{T}^7$. Since I do not know what g is (in principle there might even be several nonequivalent solutions) we will keep it as a parameter.

Let us recall the procedure for dimensional reduction to 2+1D. (see [2]). After reduction to 2+1D and Weyl-rescaling we obtain a Lagrangian of the form,

$$\begin{aligned}
L = & \int \sqrt{g} \left\{ R + \frac{2}{\Lambda^2} \partial_\mu \Lambda \partial^\mu \Lambda + \pi^2 \Lambda^4 F_{\mu\nu} F^{\mu\nu} + \frac{1}{2e^2 \Lambda^2} \partial_\mu \Phi \partial^\mu \Phi \right\} \\
& + \int \sqrt{g} \left\{ \frac{\pi^2 \Lambda^2}{e^2} H_{\mu\nu} H^{\mu\nu} + \frac{\pi^2 \Lambda^2}{e^2} \Phi^2 F_{\mu\nu} F^{\mu\nu} + \frac{2\pi^2 \Lambda^2}{e^2} \Phi H_{\mu\nu} F^{\mu\nu} \right\} \\
& + \frac{\theta}{16\pi} \int \epsilon^{\mu\nu\alpha} [H_{\mu\nu} \partial_\alpha \Phi + \Phi F_{\mu\nu} \partial_\alpha \Phi].
\end{aligned} \tag{4.1}$$

Here,

$$F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu,$$

and A_μ is the gauge field coming from the 3+1D graviton. Φ is the scalar coming from the 3+1D gauge field $\tilde{B}_\mu$, and $B_\mu \equiv \tilde{B}_\mu - \Phi A_\mu$ is the 2+1D gauge field. Λ is the radius of $\mathbf{S}^1$ (in 3+1D units). We dualize the two gauge fields A_μ and B_μ into two scalars with integral periods. After a change of variables we find a sigma-model with metric,

$$ds^2 = \frac{1}{e^4 \Lambda^4} (d\sigma_1^2 + 2\Phi d\sigma_1 d\sigma_2 + (e^2 \Lambda^2 + \Phi^2) d\sigma_2^2) + 4 \frac{d\Lambda^2}{\Lambda^2} + \frac{1}{e^2 \Lambda^2} d\Phi^2. \tag{4.2}$$

With the definition,

$$\tau = \tau_1 + i\tau_2 = \frac{2\pi i}{e^2} + \frac{\theta}{2\pi}. \tag{4.3}$$

The moduli space is subject to the identifications,

$$\begin{aligned}
\sigma_1 & \rightarrow \sigma_1 - n_3 \sigma_2 + \tau_2^{-1} n_1 - \frac{1}{2} \tau_2^{-1} \tau_1 n_3^2, \\
\sigma_2 & \rightarrow \sigma_2 + \tau_2^{-1} n_2 + \tau_2^{-1} \tau_1 n_3, \\
\Phi & \rightarrow \Phi + n_3, \\
n_1, n_2, n_3 & \in \mathbb{Z}.
\end{aligned} \tag{4.4}$$

The two coordinates $d\sigma_1 - d\sigma_2$ form a torus T^2 with complex structure,

$$\tau = ie\Lambda - \Phi,$$

and area,

$$A = \frac{e}{\Lambda^3}.$$

The target-space metric is Einstein, as required by supersymmetry [1], and satisfies,

$$R_{ab} = -\frac{3}{2} g_{ab}, \quad R = -6.$$

4.3. The symmetric space $SU(2, 1)/(SU(2) \times U(1))$

Ignoring the discrete identifications (4.4), for the moment, the moduli space and metric given by (4.2) can be identified with the symmetric Kähler manifold $SU(2, 1)/(SU(2) \times U(1))$ (see [12-14]). Let us sketch how this works. The exact details are given in appendix (D).

$SU(2, 1)$ can be realized as the subgroup of complex 3×3 matrices M with unit determinant, $SL(3, \mathbf{C})$, that preserve an indefinite form which we take to be,

$$J = \begin{pmatrix} 0 & 0 & -1 \\ 0 & 2 & 0 \\ -1 & 0 & 0 \end{pmatrix}.$$

The matrices M satisfy $M^\dagger J M = J$. We can think of the space $SU(2, 1)/(SU(2) \times U(1))$ as the space of complex directions inside $\mathbf{C}^3$ with the indefinite metric,

$$ds^2 = -dz_1 d\bar{z}_3 - dz_3 d\bar{z}_1 + 2|dz_2|^2.$$

We can parameterize the bulk of $SU(2, 1)/(SU(2) \times U(1))$ with two complex coordinates z_1, z_2 with a restriction,

$$\text{Re } z_1 > |z_2|^2. \quad (4.5)$$

The Kähler function takes the Fubini-Study form,

$$K = \log(z_1 + \bar{z}_1 - 2|z_2|^2). \quad (4.6)$$

The action of an element,

$$g \equiv \begin{pmatrix} g_{11} & g_{12} & g_{13} \\ g_{21} & g_{22} & g_{23} \\ g_{31} & g_{32} & g_{33} \end{pmatrix} \in SU(2, 1),$$

takes the form,

$$g : (z_1, z_2) \rightarrow \left(\frac{g_{11}z_1 + g_{12}z_2 + g_{13}}{g_{31}z_1 + g_{32}z_2 + g_{33}}, \frac{g_{21}z_1 + g_{22}z_2 + g_{23}}{g_{31}z_1 + g_{32}z_2 + g_{33}} \right). \quad (4.7)$$

The precise mapping between the variables $(\sigma_1, \sigma_2, \Phi, \Lambda)$ and (z_1, z_2) is written down in appendix (D). Here we will list a few simple consequences. We need to transform the identifications (4.4) to the (z_1, z_2) variables. An identification of the form,

$$(\sigma_1, \sigma_2, \Phi, \Lambda) \sim (\sigma_1 + a, \sigma_2, \Phi, \Lambda),$$

becomes,

$$(z_1, z_2) \sim (z_1 + 4ia, z_2).$$

Similarly, an identification of the form,

$$(\sigma_1, \sigma_2, \Phi, \Lambda) \sim (\sigma_1, \sigma_2 + a, \Phi, \Lambda),$$

becomes,

$$(z_1, z_2) \sim (z_1 - 2az_2 + a^2, z_2 - a),$$

and an identification of the form,

$$(\sigma_1, \sigma_2, \Phi, \Lambda) \sim (\sigma_1 - a\sigma_2, \sigma_2, \Phi + a, \Lambda),$$

becomes,

$$(z_1, z_2) \sim (z_1 + 2iaz_2 + a^2, z_2 - ia).$$

Thus, using (4.4), the classical moduli space can be taken as the space of $(z_1, z_2) \in \mathbf{C}^2$ subject to (4.5) and,

$$0 \leq \text{Im } z_2 < 1, \quad 0 \leq \text{Re } z_2 < \tau_2^{-1}, \quad 0 \leq \text{Im } z_1 < 4\tau_2^{-1}.$$

It is also important to know what is the region $\Lambda \rightarrow \infty$ in terms of (z_1, z_2) . As we show in appendix (D), this is the region,

$$\text{Re } z_1 - |z_2|^2 \rightarrow \infty.$$

The region $(\text{Re } z_1 - |z_2|^2) \rightarrow 0$ is the boundary of the moduli space and is inaccessible to classical analysis.

5. The quantum moduli space

We have seen that the moduli of dimensionally reduced 3+1D $\mathcal{N} = 2$ supergravity is given by,

$$\mathcal{M}_{cl} = \Gamma_{cl} \backslash SU(2, 1) / U(2).$$

Γ_{cl} is a discrete group generated by the shifts,

$$\begin{aligned} (z_1, z_2) &\sim (z_1 + 4i\tau_2^{-1}, z_2), \\ (z_1, z_2) &\sim (z_1 + 2\tau_2^{-1}z_2 + \tau_2^{-2}, z_2 + \tau_2^{-1}), \\ (z_1, z_2) &\sim (z_1 + 2\tau_2^{-1}\bar{\tau}z_2 + \tau_2^{-2}|\tau|^2, z_2 + \tau_2^{-1}\tau). \end{aligned} \tag{5.1}$$

As mentioned above, τ could be determined if we knew the moduli at the fixed point. After a redefinition $z_2 \rightarrow \tau_2 z_2$ and $z_1 \rightarrow |\tau_2|^2 z_1$ we find the generators of these shifts (5.1) in $SU(2, 1)$ to be,

$$g_1 = \begin{pmatrix} 1 & 0 & 4i\tau_2 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad g_2 = \begin{pmatrix} 1 & 2 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}, \quad g_3 = \begin{pmatrix} 1 & 2\bar{\tau} & |\tau|^2 \\ 0 & 1 & \tau \\ 0 & 0 & 1 \end{pmatrix}, \quad (5.2)$$

Note that,

$$g_1 = g_2^{-1} g_3^{-1} g_2 g_3$$

Note also that under $\tau \rightarrow -1/\tau$ the classical group is invariant provided we conjugate,

$$g \rightarrow P g P^{-1},$$

with,

$$P = \begin{pmatrix} -\frac{1}{\tau} & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -\bar{\tau} \end{pmatrix}.$$

What is the quantum moduli space of the theory? In principle there could be two kinds of changes from the classical limit to the full quantum result:

1. The group Γ_{cl} could be a subgroup of a larger group Γ of dualities.
2. The metric could be corrected, since we only have $\mathcal{N} = 4$ supersymmetry.

5.1. Strong coupling

The classical moduli space $SU(2, 1)/U(2)$ has a nasty infinite volume region as $\Lambda \rightarrow 0$. It is unlikely that such a singularity remains in the quantum theory. There are two conceivable ways in which this singularity is resolved. It could be that quantum corrections smooth out the moduli-space and in the quantum moduli-space only the classical region $\Lambda \rightarrow \infty$ is non-compact. The other possibility (which is my bet) is that the classical identification group Γ_{cl} is extended to a quantum duality group Γ . What are the restrictions on Γ ? Obviously, Γ must not contain any element that relates two classical-vacua, that is vacua with large Λ , that are unrelated by Γ_{cl} . As is usually the case with strong/weak dualities, the extra generators of Γ must take the weakly-coupled regime $\Lambda \rightarrow \infty$ to a strongly coupled regime of $\Lambda \sim 1$ or $\Lambda \rightarrow 0$.

As an example, let us consider the element with the matrix,

$$S = \Omega \begin{pmatrix} 0 & 0 & 1 \\ 0 & -1 & 0 \\ 1 & 0 & 0 \end{pmatrix} \Omega^{-1}.$$

For some appropriate Ω . Let us also set $\Omega = \mathbf{I}$. It is easy to check that S maps the classical region $\Lambda \rightarrow \infty$ to $\Lambda \ll 1$. For $\sigma_1 = \sigma_2 = 0$ it acts as,

$$\Phi + 2ie\Lambda \longrightarrow -\frac{1}{\Phi + 2ie\Lambda}. \quad (5.3)$$

Let Γ' be the group generated by S together with Γ_{cl} . Then,

$$\Gamma' \backslash SU(2, 1)/U(2),$$

is a good candidate for the quantum moduli space (for an appropriate Ω).

We will discuss the two possibilities, quantum corrections and an extended duality group, more in the following subsections.

5.2. Sources for the quantum corrections

The first possibility to explore is that quantum effects correct the metric. It is not completely clear to me whether mathematically such perturbation of the Quaternionic-Kähler structure is allowed. What we are looking for is a metric that preserves the behaviour at $\Lambda \rightarrow \infty$ while at the same time does not have singularities which are more nasty than ADE, and the portion of the moduli space with bounded Λ is compact. In [21] it was shown that the contribution of quantum corrections to the universal hypermultiplet metric in type-II compactifications on a Calabi-Yau can be absorbed in a redefinition of the variables. On the mathematical side, techniques for obtaining quaternionic-Kähler metrics have been developed in [44,25].² It has been shown there that the metric can be encoded in a single analytic function (denoted by $\mathcal{L}^{+4}$). We will not attempt to study the possible deformations to the metric in this paper. Below, we will list a few possible sources for quantum corrections. My bet is that these quantum corrections vanish.

From λ^{16} terms in 3+1D:

Toroidal compactifications of M-theory have calculable R^4 corrections [26-36]. They are accompanied by 16-fermion terms with a calculable coefficient [31]. In particular, M-theory on $\mathbf{T}^7$ has such a 16-fermion term, although it has not been calculated explicitly. Unless the coefficient vanishes at the particular point in moduli space in which we are working, it could correct the metric as follows. Once we compactify on $\mathbf{S}^1$, most of the fermions will not be invariant under the twist $\tilde{u}$ and will acquire a mass of order $1/\Lambda$ in 3+1D Einstein units. In 2+1D, after Weyl-rescaling the mass becomes $1/\Lambda^2$. $\mathcal{N} = 4$

² I am grateful to A. Strominger for pointing out these references.

supersymmetry in 2+1D permits 4-fermion terms. Starting with a λ^{16} term in 3+1D we get a λ^4 term if 12 fermions get a mass.

From charged particles in 3+1D:

In principle we can get terms which behave as $e^{-m\Lambda - iQ\Phi}$ from particles with mass m and charge Q under the surviving $U(1)$ gauge field in 3+1D. We can get an instanton by letting the particle have a Euclidean world-line around $\mathbf{S}^1$. There are also similar terms from the monopoles. In appendix (B) we show that such a particle is $\frac{1}{8}$ -BPS and not $\frac{1}{2}$ -BPS. Such a particle has 28 zero-modes corresponding to the broken supersymmetries. We will show in appendix (B) that 4 of the zero modes are invariant under $\tilde{u}$ while the other 24 transform with a phase and hence get a mass of the order of $\frac{1}{\Lambda}$ after the compactification on $\mathbf{S}^1$. If there were just one particle with the minimum charge we would conclude that the quantum corrections are nonzero. However, $\frac{1}{8}$ -BPS states usually have a large multiplicity. For example, to calculate the multiplicity of bound states of D2-branes on $\mathbf{T}^6$ which form a holomorphic curve inside $\mathbf{T}^6$ one has to calculate the cohomology $H^*(\mathcal{M})$ of the moduli space of such curves. The net contribution to the instanton term at lowest order would be proportional to the Euler number $\chi(\mathcal{M})$. Usually, the moduli space $\mathcal{M}$ is singular and one has to resolve it before calculating the Euler number. In the case of curves in $\mathbf{T}^4$ the Euler number turns out to be zero. It is quite likely that the net contribution is zero in our case as well.

From Kaluza-Klein monopoles:

We could also make a 2+1D instanton out of a KK monopole with respect to $\mathbf{S}^1$ which is “wrapped” over the volume of $\mathbf{T}^7$. If there were no twist, such an instanton would definitely exist and its contribution would have been proportional to $e^{-cR^2V + i\sigma_2}$. Here, V is the volume of $\mathbf{T}^7$, in 3+1D Einstein units. (This factor can be calculated from the tension of a D6-brane.) However, it is not clear to me what happens to the instanton after the twist. for large Λ , the KK monopole geometry is smooth and has a small curvature. Since our twist $\tilde{u}$ was an element of $\mathbb{Z}_6$ we can build our vacuum by modding out of M-theory on $\mathbf{T}^7 \times \widetilde{\mathbf{S}^1}$ where $\widetilde{\mathbf{S}^1}$ has a radius of 6Λ . Now we can actually construct a KK monopole solution for $\mathbf{T}^7 \times \widetilde{\mathbf{S}^1}$ and mod that solution out by $\mathbb{Z}_6$. If $\mathbb{Z}_6$ were acting freely, everything would have been fine. The problem is that $\mathbb{Z}_6$ has a fixed point at the center of the KK monopole solution. At that fixed point, the action is a combination of the geometrical $\mathbb{Z}_6$ and a non-geometrical $\tilde{u}$ and it is not clear how to study it. (See [18] for an attempt to use M(atric) theory [45] to study such cases.)

5.3. Extended group of dualities

We will now discuss in more detail the possibility that the moduli space is,

$$\Gamma \backslash SU(2,1)/U(2),$$

where $\Gamma \supset \Gamma_{cl}$ is a quantum group of dualities. We must first argue that there is a way to extend Γ_{cl} without making any unwanted classical identification and so that the dangerous region $\Lambda \rightarrow 0$ is mapped to the classical region. We will not give a general proof, but consider the following argument. Suppose we start with M-theory on $\mathbf{T}^7 \times \mathbf{S}^1$. We can embed the U-duality group $E_7(\mathbb{Z})$ of M-theory on $\mathbf{T}^7$ inside the U-duality group of M-theory on $\mathbf{T}^8$. We can then find the subgroup Γ' of the U-duality group $E_8(\mathbb{Z})$ which commutes with the element $\tilde{u}$ of the twist.³ This will be the group that preserves the form $\mathbf{T}^7 \times \mathbf{S}^1$ with $\mathbf{T}^7$ fixed at the special moduli. It is also clear that Γ' acts on $SU(2,1)/U(2)$ which is the moduli space of compactifications of the form $\mathbf{T}^7 \times \mathbf{S}^1$ with $\mathbf{T}^7$ fixed. Γ' also contains Γ_{cl} and satisfies the requirement about not making any unwanted classical identifications. Thus, if Γ' has elements which map the region $\Lambda \rightarrow 0$ to $\Lambda \rightarrow \infty$ it is a good candidate for Γ . Let us show that this is plausible. It is easier to analyze $E_8(\mathbb{Z})$ if we restrict to $\mathbf{T}^8$'s without any fluxes and with right-angles. The subgroup of $E_8(\mathbb{Z})$ which preserves these constraints turns out to be W , the Weyl group of E_8 (see [37]). Let the radii be $R_1, \dots, R_8$. Then W is generated by permutations in S_8 and by the ‘‘T-duality’’ transformation which takes the radii to,

$$V_{123}^{-2/3} R_1, V_{123}^{-2/3} R_2, V_{123}^{-2/3} R_3, V_{123}^{1/3} R_4, \dots V_{123}^{1/3} R_8, \quad V_{123} \equiv R_1 R_2 R_3.$$

Let us now perform T-duality on directions (123), then T-duality on directions (456), then T-duality on (678), then T-duality on (123), then T-duality on (145), and T-duality on (678). We finish with a permutation replacing 6 with 1, to obtain the radii,

$$\begin{aligned} r_1 &= R_1^{-1} R_2^{-2/3} R_3^{-2/3} R_4^{-1} R_5^{-1} R_6^{-1/3} R_7^{-2/3} R_8^{-2/3}, \quad r_2 = R_2^{1/3} R_3^{-2/3} R_6^{-1/3} R_7^{1/3} R_8^{1/3}, \\ r_3 &= R_2^{-2/3} R_3^{1/3} R_6^{-1/3} R_7^{1/3} R_8^{1/3}, \quad r_4 = R_2^{1/3} R_3^{1/3} R_4 R_6^{2/3} R_7^{1/3} R_8^{1/3}, \\ r_5 &= R_2^{1/3} R_3^{1/3} R_5 R_6^{2/3} R_7^{1/3} R_8^{1/3}, \quad r_6 = R_2^{-2/3} R_3^{-2/3} R_6^{-1/3} R_7^{-2/3} R_8^{-2/3}, \\ r_7 &= R_2^{1/3} R_3^{1/3} R_6^{-1/3} R_7^{1/3} R_8^{-2/3}, \quad r_8 = R_2^{1/3} R_3^{1/3} R_6^{-1/3} R_7^{-2/3} R_8^{1/3}, \end{aligned} \tag{5.4}$$

If $R_2 = R_3 = \dots = R_8 = 1$, then this transformation preserves $\mathbf{T}^7$ and takes $R_1 \rightarrow R_1^{-1}$. In principle we should analyze this not for a right-angled $\mathbf{T}^7$ but for the $\mathbf{T}^7$ in, say, the

³ I have benefited from a discussion with N. Seiberg on this point.

example of appendix (C). We believe that the conclusion will be the same, namely there exists an extra element in $E_8(\mathbb{Z})$ which preserves $\mathbf{T}^7$ and takes $\Lambda \rightarrow 0$ to $\Lambda \rightarrow \infty$.

Although, mathematically, this $\Gamma' \subset E_8(\mathbb{Z})$ has the required properties, it is not necessarily a duality in our case. The reason is that we have to make a distinction between two different ways of using $\tilde{u} \in E_7(\mathbb{Z})$ to compactify M-theory down to 2+1D. In case (i) we can mod out M-theory on $\mathbf{T}^7$ by $\tilde{u}$ already in 3+1D and then compactify the result to 2+1D. In case (ii), which is our case, we compactify from 3+1D down to 2+1D with a $\tilde{u}$ -twist. The subgroup Γ' of $E_8(\mathbb{Z})$ which preserves the moduli of $\mathbf{T}^7$ and the structure $\mathbf{T}^7 \times \mathbf{S}^1$ is indeed a duality in case (i). The disadvantage of case (i), however, is that we cannot be sure there are no new moduli from the twisted sectors. In contrast, case (ii) cannot be viewed as M-theory on $\mathbf{T}^7 \times \mathbf{S}^1$ modded out by an element of $E_8(\mathbb{Z})$. Rather, what we really mod out by is an element of $E_8(\mathbb{Z})$ compounded with a translation (by $1/6$ of the radius) along $\mathbf{S}^1$. Not all the elements of Γ' preserve the local operation of translation along $\mathbf{S}^1$. Thus, we cannot be sure that Γ' is a duality. In fact, if we were in any dimension higher than 2+1D we could most likely rule out Γ' as a duality group by studying the action on BPS states with momentum along $\mathbf{S}^1$. In 2+1D $U(1)$ charges are confined and there are no BPS states under local $U(1)$ symmetries and so the argument fails.

As an example, let us compare case (i) and case (ii) for a special case in 5+1D. We can compactify M-theory on $\mathbf{T}^4$ and then further down on $\mathbf{S}^1$ with a $\mathbb{Z}_2$ twist. If we first compactify (as in case (i)) on $\mathbf{T}^4/\mathbb{Z}_2$ we can deform to a $K3$ and we get more moduli from the twisted sectors. In case (ii) we do not get more moduli and the vector-multiplet moduli space is locally $SO(4,4)/(SO(4) \times SO(4))$, as we discussed in section (2.2) (and see also [22]). Now we can ask what happens in the limit that the radius of $\mathbf{S}^1$ is kept fixed and the volume of $\mathbf{T}^4$ shrinks to zero. In the first case we know from the duality between M-theory on $K3$ and heterotic on $\mathbf{T}^3$ that this becomes the weakly coupled heterotic theory. Now, for an appropriate choice of Wilson line along $\mathbf{S}^1$, the limit $\Lambda \rightarrow 0$ can be mapped to the T-dual heterotic string in the limit $\Lambda \rightarrow \infty$ (replacing $E_8 \times E_8$ with $SO(32)$). This comes about because the M5-branes wrapped N times on the small $K3 \times \mathbf{S}^1$ form light bound states which are then interpreted as KK states of the large T-dual dimension. So, although in this example $\Lambda \rightarrow 0$ is not dual to another point with $\Lambda \rightarrow \infty$ on the same moduli space, it is still a classical limit. Now let us see what happens in case (ii). It is likely that the M5-branes wrapped N times on the compact 5-manifold, made by the $\mathbf{T}^4$ fibered over $\mathbf{S}^1$ with the twist, do not form bound states. To see why this is reasonable, let us first compactify the $(2,0)_N$ theory of the M5-branes on a small $\mathbf{T}^4$. To a good approximation, we obtain 1+1D SYM with 16 supersymmetries. Now let us compactify

this on $\mathbf{S}^1$ with a twist. The twist “kills” 4 scalars and so we are left with 0+1D SYM with $\mathcal{N} = 8$ supersymmetries. This theory does not have a bound state [38-41]. Although the arguments for the bound state might not apply to this case because some of the fields are compact, the conclusion is likely to be correct.⁴

5.4. Phase transitions

Assuming that the moduli space is,

$$\Gamma \backslash SU(2, 1) / (SU(2) \times U(1)),$$

we can explore the singularities of the moduli space and the possible phase transitions that can occur when the moduli of the supergravity theory reach these points. The singular points are fixed points of elements of Γ . An example of such a fixed point is furnished by the element S of (5.3) that inverts $\Phi + 2ie\Lambda$ at $\sigma_1 = \sigma_2 = 0$. From (4.7) it can easily be checked that the only fixed point in the region (4.5) is at $(z_1, z_2) = (1, 0)$. This corresponds to $\Lambda = \frac{1}{2}$ and $\Phi = \sigma_1 = \sigma_2 = 0$. It follows that the local structure near that point is $\mathbb{R}^4/\mathbb{Z}_2$. This is the same structure as 2+1D $\mathcal{N} = 4$ QCD with $N_f = 2$ quarks [8,9]. The latter theory has another phase emanating from the singularity. Close to the singularity the structure of the other phase looks like $\mathbb{R}^4/\mathbb{Z}_2$ [8]. It has 3 compact parameters which parameterize $SO(3)$ and 1 non-compact parameter $0 < \rho < \infty$. We expect the same structure for small ρ in our case as well. At the singular point $\Lambda = \frac{1}{2}$ and $\Phi = \sigma_1 = \sigma_2$ the low-energy physics is described by a conformal theory whose moduli space is $\mathbb{R}^4/\mathbb{Z}_2$. At low-energies the coupling to gravity can be ignored and since QCD with $N_f = 2$ quarks (or its mirror [10]) is the only CFT which we know to possess this kind of singularity, it is natural to suspect that our supergravity theory is described at low-energies, at this particular singular point, by the same theory.

We therefore conclude that there is another phase of supergravity emanating from that point. At the other phase the original supergravity variables $\Lambda, \Phi, \sigma_1, \sigma_2$ are massive but instead we get 4 new moduli. One of them is ρ . We only know the structure of the moduli space of the other phase near $\rho = 0$. For ρ of the order of $M_p^{1/2}$, gravity mixes with the CFT again. We know from [1] that the moduli space has to be quaternionic. It is plausible that in the limit $\rho \rightarrow \infty$ (assuming the moduli-space of the other phase is non-compact) the description of the low-energy modes of the theory becomes classical again, but with a totally different description.

⁴ I have benefited from discussions with S. Sethi on this point.

6. Discussion

In the first part of this paper we have studied compactifications of M-theory with U-duality twists. We have seen that many moduli spaces from the list of [1] can be realized in this way. We have then proceeded to study a particular realization of $\mathcal{N} = 4$ 2+1D supergravity. We have given arguments for the existence of a particular twist with an isolated fixed point and which preserves $\frac{1}{4}$ of the supersymmetry. We have given the explicit form of the twist $\tilde{u}$ in $SU(8) \subset E_7(\mathbb{R})$ and we have shown that its characteristic polynomial has integral coefficients. In appendix (C) we will construct it more explicitly.

We conjectured that the moduli space of this theory is given by $\Gamma' \backslash SU(2, 1)/U(2)$ where the discrete U-duality group Γ' is an extension of the classical discrete group Γ_{cl} by a particular S-duality element. The quantum moduli space has singular points. We conjectured that the low-energy modes at the singular point in moduli space are described by the only known 2+1D conformal field-theories [10] with these types of singularity. We concluded that other phases emanate from these points. These conjectures rest on the assumption that quantum corrections do not modify the local structure of the moduli space.

In case the new phase does exist, it is an open problem whether the moduli space of the new phase is compact or not. If it is non-compact then far away from the singularity it might be described by a “classical limit” of some sort. Perhaps the theory grows more dimensions, perhaps it becomes a weakly coupled string theory, or perhaps it becomes a completely new classical limit which we have never encountered before.

Looking farther ahead, it would be interesting to understand “where” the new phases of gravity discussed in this paper “sit” with respect to the more exotic phases of gravity. These are the topological phase described in [46] and the phase with $\langle g_{\mu\nu} \rangle = 0$ suggested in [47].

Acknowledgments

I am very grateful to A.M. Polyakov, S. Sethi, E. Witten and especially N. Seiberg for discussions and to A. Strominger for correspondence.

Appendix A. Quick review of 2+1D moduli spaces

The results of [1], for more than $\mathcal{N} = 4$ supersymmetries, were summarized in the following table (extracted from table (3) of [1]).

$\mathcal{N}$	Dim	G/K
16	128	$E_{8(+8)}/SO(16)$
12	64	$E_{7(-14)}/(SO(12) \otimes SO(3))$
10	32	$E_{6(-12)}/(SO(10) \otimes SO(2))$
9	16	$F_{4(-20)}/SO(9)$
8	$8k$	$SO(8, k)/(SO(8) \otimes SO(k))$
6	$8k$	$SU(8, k)/S(U(4) \otimes U(k))$
5	$8k$	$Sp(2, k)/(Sp(2) \otimes Sp(k))$

Here $\mathcal{N}$ is the number of supersymmetries, “Dim” is the dimension of the moduli space and G/K is the local form of the moduli space.

For the case of $\mathcal{N} = 4$ the moduli space has to be quaternionic. For a 4-dimensional manifold this means that the curvature satisfies,

$$0 = R_{ijkl} + \frac{1}{2}\sqrt{g}R_{ij}{}^{mn}\epsilon_{mnkl} + g_{ik}g_{jl} - g_{il}g_{jk} + \sqrt{g}\epsilon_{mnkl}.$$

Appendix B. The fixed point of the U-duality element

In section (3), we needed a U-duality element $\tilde{u} \in E_7(\mathbb{Z})$ and an element $g \in E_{7(7)}(\mathbb{R})$ and $\tilde{k} \in SU(8)$ such that,

$$g\tilde{k}g^{-1} = \tilde{u},$$

and the eigenvalues of $\tilde{k}$ are,

$$\underbrace{(e^{\frac{i\pi}{3}}, \dots, e^{\frac{i\pi}{3}})}_{6 \text{ times}}, 1, 1). \quad (\text{B.1})$$

In this section we will make some observations on such a $\tilde{u}$, independent of a particular realization. We will start by finding the charge in M-theory on $\mathbf{T}^7$ that is fixed by $\tilde{u}$. In particular, we wish to know how much SUSY the corresponding charged particle preserves.

We start with preliminaries.

B.1. Central charge formula for M-theory on $\mathbf{T}^7$

The central charge formula for M-theory on $\mathbf{T}^7$ has been explained in [6]. The SUSY charges are in the representation,

$$(\mathbf{2}, \mathbf{8}) + (\overline{\mathbf{2}}, \overline{\mathbf{8}}),$$

of $SO(3, 1) \times SU(8)$. The charge is a vector q in the representation $\mathbf{56}$ of $E_7(\mathbb{R})$. The central charge Z is a vector in the $\mathbf{28} + \overline{\mathbf{28}}$ of $SU(8) \subset E_7(\mathbb{R})$. We pick a map T from $\mathbf{56}$ of E_7 to $\mathbf{28} + \overline{\mathbf{28}}$ of $SU(8)$. The map is required to satisfy,

$$T \circ \Omega = \Omega \circ T, \quad \Omega \in SU(8) \subset E_7(\mathbb{R})$$

Let g be a representative of the $E_7(\mathbb{R})/SU(8)$ point corresponding to the moduli. g is defined up to $g \rightarrow g\Omega$ with $\Omega \in SU(8)$. The relation between the central charge and the charge vector is,

$$\tilde{Z} = Tg^{-1}q.$$

B.2. Number of unbroken supersymmetries for BPS states

Now suppose a state has a central charge matrix of $\tilde{Z}$. What is the maximal number of supersymmetries that it can preserve? $\tilde{Z}$ can be thought of as an 8×8 antisymmetric matrix. The representation $\mathbf{28} + \overline{\mathbf{28}}$ is made from Z and Z^* . So now we can write the commutation relations,

$$\begin{aligned} \{Q_{i\alpha}, Q_{\bar{j}\dot{\beta}}\} &= \delta_{i\bar{j}} \sigma_{\alpha\dot{\beta}}^{\mu} P_{\mu}, \\ \{Q_{i\alpha}, Q_{j\beta}\} &= Z_{ij} \epsilon_{\alpha\beta}, \\ \{Q_{\bar{i}\dot{\alpha}}, Q_{\bar{j}\dot{\beta}}\} &= \overline{Z}_{\bar{i}\bar{j}} \epsilon_{\dot{\alpha}\dot{\beta}}. \end{aligned} \tag{B.2}$$

As a Matrix this is,

$$\begin{pmatrix} 0 & 0 & E\delta_{i\bar{j}} & \overline{Z}_{\bar{i}\bar{j}} \\ 0 & 0 & -Z_{ij} & E\delta_{i\bar{j}} \\ E\delta_{i\bar{j}} & Z_{ij} & 0 & 0 \\ -\overline{Z}_{\bar{i}\bar{j}} & E\delta_{i\bar{j}} & 0 & 0 \end{pmatrix}.$$

The columns and rows are in the order $1\bar{j}, 2j, 1j, 2\bar{j}$. We define the 16×16 Hermitian matrix,

$$A \equiv \begin{pmatrix} E\delta_{i\bar{j}} & \overline{Z}_{\bar{i}\bar{j}} \\ -Z_{ij} & E\delta_{i\bar{j}} \end{pmatrix} = \begin{pmatrix} E\mathbf{I} & -Z^{\dagger} \\ -Z & E\mathbf{I} \end{pmatrix}.$$

and also,

$$B = \begin{pmatrix} E\mathbf{I} & Z \\ Z^{\dagger} & E\mathbf{I} \end{pmatrix}.$$

The 32×32 matrix becomes,

$$\begin{pmatrix} 0 & A \\ B & 0 \end{pmatrix}.$$

We need to know how many zeroes A and B have together. Since A is the complex conjugate of B it is enough to count the number of zeroes of A . Thus, we are looking for,

$$0 = \begin{pmatrix} E\mathbf{I} & -Z^{\dagger} \\ -Z & E\mathbf{I} \end{pmatrix} \begin{pmatrix} u \\ v \end{pmatrix} \implies (Z^{\dagger}Z - E^2\mathbf{I})u = 0.$$

It can easily be checked that the number of supersymmetries that are preserved by the state is twice the dimension of the eigen-space of the largest eigenvalue of $Z^{\dagger}Z$. What do we know about Z ? Let g be the fixed point of the U-duality twist $\tilde{u}$. Thus,

$$\tilde{u}g = g\tilde{k}.$$

Let q be a vector of charges which is invariant under $\tilde{u}$,

$$\tilde{u}q = q.$$

It follows that,

$$\tilde{Z} = Tg^{-1}q = Tg^{-1}\tilde{u}^{-1}q = T\tilde{k}^{-1}g^{-1}q = \tilde{k}^{-1}Tg^{-1}q = \tilde{k}^{-1}\tilde{Z}.$$

This is in the representation $\mathbf{28} + \overline{\mathbf{28}}$. To bring this back to the 8×8 antisymmetric Z_{ij} we have,

$$Z = \tilde{k}Z\tilde{k}^t.$$

Taking $\tilde{k}$ as given in (B.1) we see that Z is,

$$Z = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & Z_0 \\ 0 & 0 & 0 & 0 & 0 & 0 & -Z_0 & 0 \end{pmatrix}.$$

Thus, $ZZ^\dagger$ has two eigenvalues of $|Z_0|^2$ and 6 eigenvalues of zero. We see that a BPS state with the corresponding fixed charge preserves $\frac{1}{8}$ of the supersymmetry. Out of the broken supersymmetry generators, $24 = 6 \times 4$ transform under $\tilde{u}$ with a phase and 4 generators are invariant.

Appendix C. Explicit construction of $\tilde{u}$

In section (3.2) we needed an element $\tilde{u} \in E_7(\mathbb{Z})$ and an element $g \in E_{7(7)}(\mathbb{R})$ such that $g \circ \tilde{u} \circ g^{-1}$ is equal to $\tilde{k} \in SU(8)$ in (3.10).

We will now give one explicit construction for such an element $\tilde{u} \in E_7(\mathbb{Z})$. For the construction it is convenient to view the compactification as type-II on $\mathbf{T}^6$. Let us set all the RR-fluxes to zero. The moduli space is,

$$(SO(6, 6, \mathbb{Z}) \backslash SO(6, 6, \mathbb{R}) / (SO(6) \times SO(6))) \otimes (SL(2, \mathbb{Z}) \backslash SL(2, \mathbb{R}) / SO(2)).$$

The first factor corresponds to the metric and NS-NS 2-form fluxes on $\mathbf{T}^6$ while the second part corresponds to the parameter,

$$\chi \equiv \frac{iV}{\lambda_s^2} + \tilde{C},$$

where $\tilde{C}$ is the NS-NS 6-form flux (dual of the NS-NS 2-form) on $\mathbf{T}^6$. V is the overall volume of $\mathbf{T}^6$ and λ_s is the string coupling constant. We will denote the $SO(2) \subset SL(2, \mathbb{R})$ by $U(1)_\chi$. The subgroup of the U-duality group that preserves the condition that the RR-fluxes are zero is,

$$SL(2, \mathbb{Z}) \times SO(6, 6, \mathbb{Z}).$$

At special points in the moduli space, a finite subgroup of the T-duality group $SO(6, 6, \mathbb{Z})$ becomes a symmetry. The finite subgroup can then be identified with a discrete subgroup of a cover of $SO(6) \times SO(6)$. This cover is $SU(4) \times SU(4)$. A symmetry element which can be realized as a geometrical transformation of $\mathbf{T}^6$ can be embedded in the diagonal $SO(6)$. Similarly, at special points in the moduli space, a finite subgroup of the S-duality group $SL(2, \mathbb{Z})$ becomes a symmetry. It can then be identified with a discrete subgroup of $U(1)_\chi$. Now we can embed $SU(4) \times SU(4) \times U(1)_\chi$ inside $SU(8)$ which we have used to write down (3.10). Under this embedding,

$$\mathbf{8} = \mathbf{4}_+ + \mathbf{4}_-.$$

Now let us take a special $\mathbf{T}^6$ of the form $\mathbf{T}^2_a \times \mathbf{T}^2_b \times \mathbf{T}^2_c$. To make the discussion clearer, we have added subscripts to the $\mathbf{T}^2$'s. Let us denote the complex structure of each $\mathbf{T}^2$ by τ (with an appropriate subscript) and the combination $\frac{B}{2\pi} + iA$ by ρ . Here B is the NS-NS 2-form and A is the area. Type-II on $\mathbf{T}^2$ has a moduli space of,

$$(SL(2, \mathbb{Z}) \backslash SL(2, \mathbb{R}) / U(1)_\rho) \times (SL(2, \mathbb{Z}) \backslash SL(2, \mathbb{R}) / U(1)_\tau)$$

corresponding to the pair (ρ, τ) . If we write the moduli space in the form,

$$SO(2, 2, \mathbb{Z}) \backslash SO(2, 2, \mathbb{R}) / (SO(2)_1 \times SO(2)_2),$$

then $U(1)_\tau$ is the diagonal combination of $SO(2)_1$ and $SO(2)_2$ while $U(1)_\rho$ is the combination with $SO(2)_2$ inverted. As before, at special points in the moduli space we can identify the finite subgroup of the U-duality group which preserves the point and the structure $\mathbf{T}^2_c \times \mathbf{T}^2_b \times \mathbf{T}^2_a$ with a discrete subgroup of,

$$U(1)_{\rho_a} \times U(1)_{\tau_a} \times U(1)_{\rho_b} \times U(1)_{\tau_b} \times U(1)_{\rho_c} \times U(1)_{\tau_c} \times U(1)_\lambda \subset SU(8). \quad (\text{C.1})$$

The representation $\mathbf{8}$ decomposes as,

$$\begin{aligned} & (+ + + + + + +) \oplus (+ + - - - - +) \oplus (- - + + - - +) \oplus (- - - - + + +) \oplus \\ & (+ - + - + - -) \oplus (+ - - + - + -) \oplus (- + + - - + -) \oplus (- + - + + - -) \end{aligned} \quad (\text{C.2})$$

Now we take $\mathbf{T}_a^2$ to have $\tau_a = \rho_a = e^{-\frac{2\pi i}{3}}$ and $\mathbf{T}_b^2$ and $\mathbf{T}_c^2$ to have,

$$\tau_b = \tau_c = \rho_b = \rho_c = i.$$

We also take,

$$\chi = e^{-\frac{2\pi i}{3}}.$$

The U-duality element $\tilde{u}$ is a combination of,

$$\begin{aligned} \tau_b &\rightarrow -\frac{1}{\tau_b}, \quad \rho_b \rightarrow -\frac{1}{\rho_b}, & \tau_c &\rightarrow -\frac{1}{\tau_c}, \quad \rho_c \rightarrow -\frac{1}{\rho_c}, \\ \tau_a &\rightarrow -1 - \frac{1}{\tau_a}, \quad \rho_a \rightarrow -1 - \frac{1}{\rho_a}, \end{aligned} \tag{C.3}$$

and we top it with an S-duality transformation,

$$\chi \rightarrow -1 - \frac{1}{\chi}.$$

This corresponds to the following elements in (C.1):

$$(e^{\frac{-\pi i}{3}}, e^{\frac{-\pi i}{3}}, e^{\frac{\pi i}{4}}, e^{\frac{\pi i}{4}}, e^{\frac{\pi i}{4}}, e^{\frac{\pi i}{4}}, e^{\frac{-\pi i}{3}}).$$

Note that since all of these transformations can be U-conjugated to rotations of a $\mathbf{T}^2$ and since the $U(1)$ charges are measured with respect to the fermions, the phases are half the rotation angle. (The overall $(-)^F$ is a symmetry of M-theory.) Using (C.2) we calculate the eigenvalues of $\tilde{k}$ in $\mathbf{8}$ of $SU(8)$ to be,

$$(1, 1, e^{\frac{\pi i}{3}}, e^{\frac{\pi i}{3}}, e^{\frac{\pi i}{3}}, e^{\frac{\pi i}{3}}, e^{\frac{\pi i}{3}}, e^{\frac{\pi i}{3}}).$$

C.1. The invariant charge

We can now calculate the invariant charge corresponding to the $\frac{1}{8}$ -BPS particle (see appendix B). The representation of the charges is the $\mathbf{56}$ of $E_7(\mathbb{Z})$. Under,

$$SO(6, 6, \mathbb{Z}) \times SL(2, \mathbb{Z}),$$

it decomposes as,

$$\mathbf{56} = (\mathbf{12}, \mathbf{2}) + (\mathbf{32}, \mathbf{1}),$$

(see [6]). Under $SO(2, 2)_a \times SO(2, 2)_b \times SO(2, 2)_c \times SL(2)_\chi$ it decomposes as,

$$\begin{aligned} \mathbf{56} = & (\mathbf{4}, \mathbf{1}, \mathbf{1}, \mathbf{2}) + (\mathbf{1}, \mathbf{4}, \mathbf{1}, \mathbf{2}) + (\mathbf{1}, \mathbf{1}, \mathbf{4}, \mathbf{2}) + \\ & (\mathbf{2}, \mathbf{2}, \mathbf{2}, \mathbf{1}) + (\mathbf{2}, \mathbf{2}', \mathbf{2}', \mathbf{1}) + (\mathbf{2}', \mathbf{2}, \mathbf{2}', \mathbf{1}) + (\mathbf{2}', \mathbf{2}', \mathbf{2}, \mathbf{1}) \end{aligned} \tag{C.4}$$

Under $U(1)_{\rho_a} \times U(1)_{\tau_a}$ of (C.1), $\mathbf{4}$, $\mathbf{2}$ and $\mathbf{2}'$ of $SO(2, 2)_a$ decompose as,

$$\begin{aligned}\mathbf{4} &= (+1, +1) \oplus (+1, -1) \oplus (-1, +1) \oplus (-1, -1), \\ \mathbf{2} &= (+1, 0) \oplus (-1, 0), \\ \mathbf{2}' &= (0, +1) \oplus (0, -1),\end{aligned}\tag{C.5}$$

(If we replace type-IIA with type-IIB, we have to replace ρ_a with τ_a .) We see that two vectors out of, $(\mathbf{4}, \mathbf{1}, \mathbf{1}, \mathbf{2})$, are invariant under $\tilde{u}$. The corresponding BPS states are combinations of strings wrapped on $\mathbf{T}^2_a$ and KK states along the sides of $\mathbf{T}^2_a$ together with their duals which are NS5-branes and KK monopoles.

Appendix D. The $SU(2, 1)/(SU(2) \times U(1))$ moduli space

We have seen that the target-space with the metric,

$$ds^2 = \frac{1}{e^4 \Lambda^4} (d\sigma_1^2 + 2\Phi d\sigma_1 d\sigma_2 + (e^2 \Lambda^2 + \Phi^2) d\sigma_2^2) + 4 \frac{d\Lambda^2}{\Lambda^2} + \frac{1}{e^2 \Lambda^2} d\Phi^2, \tag{D.1}$$

is equivalent to the space $SU(2, 1)/(SU(2) \times U(1))$ and the latter can be described in terms two complex coordinates (z_1, z_2) , with the action of an $SU(2, 1)$ matrix,

$$g = \begin{pmatrix} g_{11} & g_{12} & g_{13} \\ g_{21} & g_{22} & g_{23} \\ g_{31} & g_{32} & g_{33} \end{pmatrix},$$

given by,

$$g : (z_1, z_2) \rightarrow \left(\frac{g_{11}z_1 + g_{12}z_2 + g_{13}}{g_{31}z_1 + g_{32}z_2 + g_{33}}, \frac{g_{21}z_1 + g_{22}z_2 + g_{23}}{g_{31}z_1 + g_{32}z_2 + g_{33}} \right).$$

The purpose of this appendix is to describe the precise mapping.

D.1. Killing vectors

The metric (D.1) possesses a group of isometries which can be described by the following infinitesimal transformation laws,

$$\begin{aligned}\delta\sigma_1 &= \epsilon_1 \left\{ \frac{3}{2} \Phi^2 \sigma_2 - \frac{1}{2} \sigma_2^3 \right\} + \epsilon_2 \left\{ \frac{3}{8} \Phi^4 + 2e^2 \Lambda^2 \Phi^2 + 2e^4 \Lambda^4 - 2\sigma_1^2 + \frac{3}{4} \Phi^2 \sigma_2^2 - \frac{1}{8} \sigma_2^4 \right\} \\ &\quad - 2\epsilon_3 \sigma_1 - \epsilon_4 \{ \Phi^3 + 2e^2 \Lambda^2 \Phi + 2\sigma_1 \sigma_2 \} - \epsilon_5 \sigma_2 + \epsilon_6 \left\{ \frac{1}{2} \Phi^2 - \frac{1}{2} \sigma_2^2 \right\} + \epsilon_8, \\ \delta\sigma_2 &= -\epsilon_1 \{ 2\sigma_1 + 3\Phi \sigma_2 \} - \epsilon_2 \left\{ \frac{1}{2} \Phi^3 + 2e^2 \Lambda^2 \Phi + 2\sigma_1 \sigma_2 + \frac{3}{2} \Phi \sigma_2^2 \right\} \\ &\quad - \epsilon_3 \sigma_2 + \epsilon_4 \left\{ \frac{3}{2} \Phi^2 + 2e^2 \Lambda^2 - \frac{1}{2} \sigma_2^2 \right\} - \epsilon_6 \Phi + \epsilon_7 \\ \delta\Phi &= \epsilon_1 \left\{ 2e^2 \Lambda^2 - \frac{1}{2} \Phi^2 + \frac{3}{2} \sigma_2^2 \right\} + \epsilon_2 \left\{ 2e^2 \Lambda^2 \sigma_2 - \frac{1}{2} \Phi^2 \sigma_2 - 2\Phi \sigma_1 + \frac{1}{2} \sigma_2^3 \right\} \\ &\quad - \epsilon_3 \Phi + \epsilon_4 \{ 2\sigma_1 - \Phi \sigma_2 \} + \epsilon_5 + \epsilon_6 \sigma_2, \\ \delta\Lambda &= -\epsilon_1 \Phi \Lambda - \epsilon_2 \{ 2\Lambda \sigma_1 + \Phi \Lambda \sigma_2 \} - \epsilon_3 \Lambda - \epsilon_4 \Lambda \sigma_2.\end{aligned}\tag{D.2}$$

Here $\epsilon_1, \dots, \epsilon_8$ are arbitrary.

Writing,

$$\delta \equiv \sum_{i=1}^8 \epsilon_i \delta_i,$$

we can calculate the commutation relations,

$$\begin{aligned} [\delta_1, \delta_2] &= 0, & [\delta_1, \delta_3] &= -\delta_1, & [\delta_1, \delta_4] &= 2\delta_2, & [\delta_1, \delta_5] &= \delta_3, \\ [\delta_1, \delta_6] &= \delta_4, & [\delta_1, \delta_7] &= 3\delta_6, & [\delta_1, \delta_8] &= -2\delta_7, & [\delta_2, \delta_3] &= -2\delta_2, \\ [\delta_2, \delta_4] &= 0, & [\delta_2, \delta_5] &= -\delta_4, & [\delta_2, \delta_6] &= 0, & [\delta_2, \delta_7] &= \delta_1, \\ [\delta_2, \delta_8] &= 2\delta_3, & [\delta_3, \delta_4] &= \delta_4, & [\delta_3, \delta_5] &= -\delta_5, & [\delta_3, \delta_6] &= 0, \end{aligned} \quad (\text{D.3})$$

$$\begin{aligned} [\delta_3, \delta_7] &= -\delta_7, & [\delta_3, \delta_8] &= -2\delta_8, & [\delta_4, \delta_5] &= -3\delta_6, & [\delta_4, \delta_6] &= -\delta_1, \\ [\delta_4, \delta_7] &= \delta_3, & [\delta_4, \delta_8] &= 2\delta_5, & [\delta_5, \delta_6] &= \delta_7, & [\delta_5, \delta_7] &= -\delta_8, \\ [\delta_5, \delta_8] &= 0, & [\delta_6, \delta_7] &= \delta_5, & [\delta_6, \delta_8] &= 0, & [\delta_7, \delta_8] &= 0, \end{aligned} \quad (\text{D.4})$$

These generators form the Lie algebra of $SU(2, 1)$. We can represent the algebra as the set of 3×3 matrices A which satisfy,

$$A^\dagger J = -JA,$$

where,

$$J = \begin{pmatrix} 0 & 0 & -1 \\ 0 & 2 & 0 \\ -1 & 0 & 0 \end{pmatrix}.$$

The generators can be represented as,

$$\begin{aligned} \delta_1 &= \begin{pmatrix} 0 & 0 & 0 \\ -\frac{1}{2}i & 0 & 0 \\ 0 & i & 0 \end{pmatrix}, & \delta_2 &= \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ -\frac{1}{2}i & 0 & 0 \end{pmatrix}, \\ \delta_3 &= \begin{pmatrix} -1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}, & \delta_4 &= \begin{pmatrix} 0 & 0 & 0 \\ -\frac{1}{2} & 0 & 0 \\ 0 & -1 & 0 \end{pmatrix}, \end{aligned} \quad (\text{D.5})$$

$$\begin{aligned} \delta_5 &= \begin{pmatrix} 0 & 2i & 0 \\ 0 & 0 & -i \\ 0 & 0 & 0 \end{pmatrix}, & \delta_6 &= \begin{pmatrix} -\frac{i}{3} & 0 & 0 \\ 0 & \frac{2i}{3} & 0 \\ 0 & 0 & -\frac{i}{3} \end{pmatrix}, \\ \delta_7 &= \begin{pmatrix} 0 & -2 & 0 \\ 0 & 0 & -1 \\ 0 & 0 & 0 \end{pmatrix}, & \delta_8 &= \begin{pmatrix} 0 & 0 & 4i \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}. \end{aligned} \quad (\text{D.6})$$

D.2. Integrated forms

We will now write down some group element actions of the form $e^{t\delta_i}$. It is easy to see that,

$$\begin{aligned} e^{t\delta_8}(\sigma_1, \sigma_2, \Phi, \Lambda) &= (\sigma_1 + t, \sigma_2, \Phi, \Lambda), \\ e^{t\delta_7}(\sigma_1, \sigma_2, \Phi, \Lambda) &= (\sigma_1, \sigma_2 + t, \Phi, \Lambda), \\ e^{t\delta_5}(\sigma_1, \sigma_2, \Phi, \Lambda) &= (\sigma_1 - t\sigma_2, \sigma_2, \Phi + t, \Lambda), \\ e^{t\delta_3}(\sigma_1, \sigma_2, \Phi, \Lambda) &= (e^{-2t}\sigma_1, e^{-t}\sigma_2, e^{-t}\Phi, e^{-t}\Lambda). \end{aligned} \tag{D.7}$$

It is also easy to integrate,

$$\begin{aligned} e^{t\delta_6}\sigma_1 &= \sigma_1 + \frac{1}{4}(\Phi^2 - \sigma_2^2)\sin 2t + \frac{1}{2}\Phi\sigma_2(1 - \cos 2t), \\ e^{t\delta_6}\sigma_2 &= \sigma_2 \cos t - \Phi \sin t, \\ e^{t\delta_6}\Phi &= \Phi \cos t + \sigma_2 \sin t, \\ e^{t\delta_6}\Lambda &= \Lambda. \end{aligned} \tag{D.8}$$

Now let us check the flow generated by $\delta_1 - \frac{1}{2}C^2\delta_5$.

$$\begin{aligned} \frac{d\sigma_1}{dt} &= \frac{3}{2}\Phi^2\sigma_2 - \frac{1}{2}\sigma_2^3 + \frac{1}{2}C^2\sigma_2, \\ \frac{d\sigma_2}{dt} &= -2\sigma_1 - 3\Phi\sigma_2, \\ \frac{d\Phi}{dt} &= 2e^2\Lambda^2 - \frac{1}{2}\Phi^2 + \frac{3}{2}\sigma_2^2 - \frac{1}{2}C^2, \\ \frac{d\Lambda}{dt} &= -\Phi\Lambda. \end{aligned} \tag{D.9}$$

Let us solve it for $\sigma_1 = \sigma_2 = 0$. We then define $W = \Phi + 2ie\Lambda$. The equations are,

$$\frac{dW}{dt} = -\frac{1}{2}W^2 - \frac{1}{2}C^2.$$

Thus,

$$W = \frac{W_0 - C \tan \frac{1}{2}Ct}{1 + \frac{W_0}{C} \tan \frac{1}{2}Ct}.$$

For $Ct = \pi$ this becomes the transformation $W_0 \rightarrow -C^2/W_0$ which we will interpret as a strong/weak duality. For $C = 1$ The matrix representation gives,

$$e^{\pi(\delta_1 - \frac{1}{2}\delta_5)} = \begin{pmatrix} 0 & 0 & 1 \\ 0 & -1 & 0 \\ 1 & 0 & 0 \end{pmatrix}.$$

D.3. Equations for the change of variables

We can now write down the equations for the change of variables from the $(\sigma_1, \sigma_2, \Phi, \Lambda)$ variables to (z_1, z_2) . To do this we will use a combination of δ_3 and $\delta_5, \delta_7, \delta_8$ to move from one point on the manifold to another,

$$e^{t\delta_3} e^{b\delta_5} e^{a_1\delta_8 + a_2\delta_7}(\sigma_1, \sigma_2, \Phi, \Lambda) = (e^{-2t}\sigma_1 - be^{-t}\sigma_2 + a_1, e^{-t}\sigma_2 + a_2, e^{-t}\Phi + b, e^{-t}\Lambda). \quad (\text{D.10})$$

We now take,

$$\begin{aligned} t &= \log \frac{\Lambda}{\tilde{\Lambda}}, \\ b &= \tilde{\Phi} - \frac{\Lambda}{\tilde{\Lambda}}\Phi, \\ a_2 &= \tilde{\sigma}_2 - \frac{\Lambda}{\tilde{\Lambda}}\sigma_2, \\ a_1 &= \tilde{\sigma}_1 - \frac{\Lambda^2}{\tilde{\Lambda}^2}\sigma_1 + \frac{\Lambda}{\tilde{\Lambda}}b\sigma_2. \end{aligned} \quad (\text{D.11})$$

The rescaling transformation $e^{t\delta_3}$ also makes it obvious that the classical regime $\Lambda \rightarrow \infty$ is given by

$$\text{Re } z_1 - |z_2|^2 \rightarrow \infty.$$

Note that neither of $\delta_5, \delta_7, \delta_8$ change the value of $\text{Re } z_1 - |z_2|^2$, as it should be.

Appendix E. On the structure of Γ for $\tau = 2i$

We have defined $\Gamma' \subset SU(2, 1)$ as the subgroup generated by,

$$g_2 = \begin{pmatrix} 1 & 2 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}, \quad g_3 = \begin{pmatrix} 1 & 2\bar{\tau} & |\tau|^2 \\ 0 & 1 & \tau \\ 0 & 0 & 1 \end{pmatrix}, \quad (\text{E.1})$$

and,

$$S = \begin{pmatrix} 0 & 0 & 1 \\ 0 & -1 & 0 \\ 1 & 0 & 0 \end{pmatrix}.$$

In this appendix we will prove that, for the special case of $\tau = 2i$, adding S does not cause any unwanted identifications in the classical region. I do not know how to generalize the proof to other τ 's. Perhaps, embedding Γ' in $E_8(\mathbb{Z})$ as the subgroup that preserves $\tilde{u}$ would be a good tactic. Translated to matrices, this statement means that there is no element $g \in \Gamma - \Gamma_{cl}$ such that the vector,

$$\begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}$$

is an eigen-vector of g .

E.1. The boundary of $SU(2, 1)/(SU(2) \times U(1))$

To understand Γ , it is helpful to analyze its action on the boundary of $SU(2, 1)/(SU(2) \times U(1))$. This boundary is given by $\text{Re } z_1 = |z_2|^2$ (equation (4.5)). Adding a point at infinity, the boundary is easily seen to be $\mathbf{S}^3$ (this is easier to see in a diagonal metric of signature $(1, 1, -1)$).

The classical region is given by $\text{Re } z_1 \gg |z_2|^2$ but it is easy to see that this is just the point at infinity that was added. Thus, in order to prove that Γ does not make extra identifications in the classical region, we need to analyze the action of Γ on the boundary.

E.2. The structure of Γ

We set $\tau = 2i$ and find,

$$g_1 = g_3^{-1} g_2^{-1} g_3 g_2 = \begin{pmatrix} 1 & 0 & 8i \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

Now let us introduce,

$$g(k, l, m) \equiv g_1^m g_2^k g_3^l = \begin{pmatrix} 1 & 2(k - 2il) & (k^2 + 4l^2) + 4i(kl + 2m) \\ 0 & 1 & (k + 2il) \\ 0 & 0 & 1 \end{pmatrix}.$$

We have the rule,

$$g(k_1, l_1, m_1)g(k_2, l_2, m_2) = g(k_1 + k_2, l_1 + l_2, m_1 + m_2 + k_2 l_1).$$

A generic term in Γ is of one of the three forms,

$$\begin{aligned} h_1 &= Sg(k_1, l_1, m_1)Sg(k_2, l_2, m_2)Sg(k_3, l_3, m_3) \cdots Sg(k_r, l_r, m_r), \\ h_2 &= g(k_1, l_1, m_1)Sg(k_2, l_2, m_2)Sg(k_3, l_3, m_3) \cdots Sg(k_r, l_r, m_r), \\ h_3 &= g(k_1, l_1, m_1)Sg(k_2, l_2, m_2)Sg(k_3, l_3, m_3) \cdots Sg(k_r, l_r, m_r)S. \\ h_4 &= Sg(k_1, l_1, m_1)Sg(k_2, l_2, m_2)Sg(k_3, l_3, m_3) \cdots Sg(k_r, l_r, m_r)S. \end{aligned} \tag{E.2}$$

We can also require that no $g(0, \pm 1, 0)$ ever appear between two S 's because we could then reduce the number of S 's (denoted by r) according to,

$$Sg(\pm 1, 0, 0)S = g(\mp 1, 0, 0)Sg(\mp 1, 0, 0).$$

Now we need a lemma.

Lemma:

Let g be given by,

$$g = Sg(k_1, l_1, m_1) Sg(k_2, l_2, m_2) Sg(k_3, l_3, m_3) \cdots Sg(k_r, l_r, m_r), \quad (\text{E.3})$$

Such that for all j ,

$$(k_j, l_j, m_j) \neq (\pm 1, 0, 0),$$

Let,

$$x = (z_1 = |z_2|^2 + iw, z_2), \quad z_2 \in \mathbf{C}, \quad w \in \mathbb{R},$$

be a point on the boundary that satisfies

$$0 < |z_2| < 1, \quad -\frac{1}{2} < w < \frac{1}{2}, \quad (\text{E.4})$$

Let x' be the result of the action of g on x (which is still a point on the boundary),

$$x' \equiv gx = (|z'_2|^2 + iw', z'_2).$$

then $0 < |z'_2| < 1$ and $|w'| < \frac{1}{2}$.

Proof:

It is sufficient to check for,

$$g = Sg(k, l, m).$$

Let us denote,

$$\mu = k + 2il, \quad \zeta = 4kl + 8m.$$

From (4.7) we find,

$$\begin{aligned} z'_2 &= -\frac{\mu + z_2}{|z_2|^2 + 2\bar{\mu}z_2 + |\mu|^2 + i(w + \zeta)}, \\ w' &= \frac{w + \zeta + 2\text{Im}(\bar{\mu}z_2)}{||z_2|^2 + 2\bar{\mu}z_2 + |\mu|^2 + i(w + \zeta)|^2}, \end{aligned} \quad (\text{E.5})$$

First, let us assume that $|\mu| \geq 2$.

$$||z_2|^2 + 2\bar{\mu}z_2 + |\mu|^2 + i(w + \zeta)| \geq ||\mu|^2 + 2\text{Re}(\bar{\mu}z_2) + |z_2|^2| = |\mu + z_2|^2. \quad (\text{E.6})$$

Thus,

$$|z'_2| \leq \frac{1}{|z_2 + \mu|} \leq \frac{1}{|\mu| - |z_2|} < 1.$$

Also let us denote by,

$$x = w + \zeta + 2\text{Im}(\bar{\mu}z_2).$$

Then,

$$|w'| = \frac{|x|}{||z_2|^2 + 2\text{Re} \bar{\mu}z_2 + |\mu|^2 + |x|^2} = \frac{|x|}{|z_2 + \mu|^2 + |x|^2} < \frac{|x|}{1 + |x|^2} \leq \frac{1}{2}, \quad (\text{E.7})$$

since $|z_2 + \mu| > 1$. To complete the proof, we need to check the case $\mu = 0$. In this case we know that $|\zeta| \geq 8$ and the same conclusion follows.

Now we can prove that all the terms in (E.2) are different from unity. For this purpose we start with the point $(z_1 = \infty, z_2)$ such that z_2 is finite. This is the classical region of the moduli space. Acting either with S or $Sg(k_r, l_r, m_r)$ will turn that point into $(0, 0)$. The next $Sg(k, l, m)$ which will act on it, will bring it into the region (E.4). The successive $Sg(k_i, l_i, m_i)$ will not be able to take it out of (E.4). If we have an extra final S (as in h_2 and h_3 of (E.2)) it will not bring it to the point at infinity because $|z_2| > 0$.

References

- [1] B. de Wit, A.K. Tollsten and H. Nicolai, “Locally Supersymmetric $D = 3$ Nonlinear Sigma-Models,” Nucl. Phys. **B392** (1993) 3, *hep-th/9208074*
- [2] N. Marcus and J.H. Schwarz, “Inifinite Symmetry Algebras of Extended Supergravity Theories,” Nucl. Phys. **B243** (1984) 243
- [3] C. M. Hull and P.K. Townsend, “Unity Of Superstring Dualities,” Nucl. Phys. **B438** (95) 109, *hep-th/9410167*
- [4] A. Sen, “Strong Weak Coupling Duality in Three Dimensional String Theory,” Nucl. Phys. **B434** (1995) 179, *hep-th/9408083*
- [5] N. Seiberg and E. Witten, “Monopoles, Duality and Chiral Symmetry Breaking in $N = 2$ Supersymmetric QCD,” Nucl. Phys. **B431** (1994) 484–550, *hep-th/9408099*
- [6] E. Witten, “String Theory Dynamics in Various Dimensions,” Nucl. Phys. **B443** (95) 85, *hep-th/9503124*
- [7] P. C. Argyres, M. R. Plesser, N. Seiberg, E. Witten, “New $N = 2$ Superconformal Field Theories In Four Dimensions,” Nucl. Phys. **B461** (96) 71, *hep-th/9511154*
- [8] N. Seiberg, “IR Dynamics on Branes and Space-Time Geometry,” Phys. Lett. **384B** (1996) 81–85, *hep-th/9606017*
- [9] N. Seiberg and E. Witten, “Gauge Dynamics And Compactification To Three Dimensions,” *hep-th/9607163*
- [10] K. Intriligator and N. Seiberg, “Mirror Symmetry in Three Dimensional Gauge Theories,” Phys. Lett. **387B** (1996) 513, *hep-th/9607207*
- [11] O.J. Ganor, D.R. Morrison and N. Seiberg, “Branes, Calabi-Yau Spaces, and Toroidal Compactification of the $N = 1$ Six-Dimensional E_8 Theory,” Nucl. Phys. **B487** (1997) 93, *hep-th/9610251*
- [12] S.Cecotti, S.Ferrara and L.Girardello, “Geometry of Type-II Superstrings and the Moduli of Superconformal Field Theories,” Int. Jour. Mod. Phys. **A4** (1989) 2475
- [13] S. Ferrara and S. Sabharwal, “Quaternionic Manifolds for type-II Superstring Vacua of Calabi-Yau Spaces,” Nucl. Phys. **B332** (1990) 317
- [14] B. de Wit, Grisaru, E. Rabinovici and H. Nicolai, Phys. Lett. **286B** (1992) 78
- [15] A. Sen and C. Vafa, “Dual Pairs of Type-II String Compactifications,” Nucl. Phys. **B455** (1995) 165, *hep-th/9508064*
- [16] K. Dasgupta and S. Mukhi, “F-Theory at Constant Coupling,” Phys. Lett. **385B** (1996) 125, *hep-th/9606044*
- [17] A. Kumar and C. Vafa, “U-Manifolds,” Phys. Lett. **396B** (1997) 85-90, *hep-th/9611007*
- [18] M. Dine and E. Silverstein, “New M-theory Backgrounds with Frozen Moduli,” *hep-th/9712166*

- [19] S. Kachru and E. Silverstein, “Selfdual Nonsupersymmetric Type-II String Compactifications,” *hep-th/9808056*
- [20] K. Becker, M. Becker and A. Strominger, unpublished.
- [21] A. Strominger, “Loop Corrections to the Universal Hypermultiplet,” *Phys. Lett.* **421B** (1998) 139, *hep-th/9706195*
- [22] A. Dabholkar and J.A. Harvey, “String Islands,” *hep-th/9809122*
- [23] S. Kachru and C. Vafa, “Exact Results For $N = 2$ Compactifications Of Heterotic Strings,” *Nucl. Phys.* **B450** (95) 69, *hep-th/9505105*
- [24] O.J. Ganor, “A Note On Zeroes of Superpotentials in F-Theory,” *Nucl. Phys.* **B499** (1997) 55, *hep-th/9612077*
- [25] E. Ivanov and G. Valent, “Quaternionic Taub-NUT from the harmonic space approach,” *hep-th/9809108*
- [26] M.B. Green, M. Gutperle and P. Vanhove, “One loop in eleven dimensions,” *hep-th/9706175*
- [27] E. Kiritsis and B. Pioline, “On R^4 threshold corrections in IIB string theory and (p, q) string instantons,” *hep-th/9707018*
- [28] I. Antoniadis, B. Pioline and T.R. Taylor, “Calculable $e^{-1/\lambda}$ Effects,” *hep-th/9707222*
- [29] A. Gregori, E. Kiritsis, C. Kounnas, N.A. Obers, P.M. Petropoulos, and B. Pioline, “ R^2 Corrections and Non-perturbative Dualities of $N = 4$ String ground states,” *Nucl. Phys.* **B510** (1998) 423–476, *hep-th/9708062*
- [30] E. Kiritsis and B. Pioline, “U-duality and D-brane Combinatorics,” *hep-th/9710078*
- [31] M.B. Green and M. Gutperle, “ λ^{16} and related terms in M-theory on T^2 ,” *hep-th/9710151*
- [32] M.B. Green and M. Gutperle, “D-particle bound states and the D-instanton measure,” *hep-th/9711107*
- [33] N.A. Obers, B. Pioline and E. Rabinovici, “M-Theory and U-duality on T^d with Gauge Backgrounds,” *hep-th/9712084*
- [34] B. Pioline, “D-effects in Toroidally Compactified Type II String Theory,” *hep-th/9712155*
- [35] M. Gutperle, “Aspects of D-Instantons,” *hep-th/9712156*
- [36] M.B. Green and S. Sethi, “Supersymmetry Constraints on Type-IIB Supergravity,” *hep-th/9808061*
- [37] S. Elizur, A. Giveon, D. Kutasov and E. Rabinovici, “Algebraic Aspects of Matrix Theory on T^d ,” *hep-th/9707217*
- [38] S. Sethi and M. Stern, “D-Brane Bound States Redux,” *hep-th/9705046*
- [39] P. Yi, “Witten Index and Threshold Bound States of D-Branes,” *hep-th/9704098*
- [40] M. Porrati and A. Rozenberg, “Bound States at Threshold in Supersymmetric Quantum Mechanics,” *hep-th/9708119*

- [41] V.A. Kazakov, I.K. Kostov and N.A. Nekrasov, “D-particles, Matrix Integrals and KP hierarchy,” *hep-th/9810035*
- [42] E. Cremmer and B. Julia, “The $SO(8)$ Supergravity,” Nucl. Phys. **B159** (79) 141
- [43] E. Cremmer, B. Julia and J. Scherk, “Supergravity Theory in Eleven-Dimensions,” Phys. Lett. **76B** (78) 409
- [44] A. Galperin, E. Ivanov and O. Ogievetskii, *hep-th/9212155*, Ann. Phys. **230** (1994) 201
- [45] T. Banks, W. Fischler, S.H. Shenker and L. Susskind, “M Theory As A Matrix Model: A Conjecture,” *hep-th/9610043*, Phys. Rev. **D55** (1997) 5112-5128
- [46] E. Witten, “Topological Gravity,” Phys. Lett. **206B** (1988) 601
- [47] A.M. Polyakov, “A Few Projects In String Theory,” *hep-th/9304146*, Published in Les Houches Sum.Sch.1992:0783-804 (QC178:H6:1992)